

Examples of our experience in supporting clinical research or questionnaire development
(published papers with our employee's names or our name in author, acknowledgement or text)

Title/Journal	Listed as/in
Use of a healthcare claims database for post-marketing safety assessments of eribulin in Japan: A comparative assessment with a prospective post-marketing surveillance study. <i>Drugs Real World Outcomes</i> . 2019; 6(1): 27-35.	Author
Secular trend in the incidence of Japanese employees with thyroid cancer undergoing thyroidectomy from 2005 to 2014: A retrospective descriptive study using an employment-based insurance claims database. <i>J of thyroid cancer</i> . 2019; 1: 5-15.	-
Ultra-high-dose methylcobalamin in amyotrophic lateral sclerosis: A long-term phase II/III randomised controlled study. <i>J Neurol Neurosurg Psychiatry</i> . Pulished first online Jan 13, 2019.	Acknowledgment
Differences in the incidence of postoperative pneumonia after percutaneous endoscopic gastrostomy between liquid and semi-solid nutrient administration. <i>Eur J Clin Nutr</i> . 2019; 73(2): 250-257.	Acknowledgment and text (methods)
Assessment of risk factors for non-specific chronic disabling low back pain in Japanese workers—findings from the CUPID (Cultural and Psychosocial Influences on Disability) study. <i>Industrial Health</i> . Published first online Oct 19, 2018	Author
Greater burden of chronic comorbidities and co-medications among people living with HIV versus people without HIV in Japan: A hospital claims database study. <i>J Infect Chemother</i> . 2019; 25(2): 89-95.	Acknowledgment
Chemotherapy-induced peripheral neuropathy in breast cancer patients treated with eribulin: interim data from a post-marketing observational study. <i>Breast Cancer</i> . 2019; 26(2): 235-243.	Acknowledgment
Pondering the “psychogenic pain”: Proposal for using the term “cognitively perceived pain”. <i>Pain Research</i> . 2018; 33(3): 183-192.	Disclosure
Healthcare burden of recurrent <i>Clostridioides difficile</i> infection in Japan: A retrospective database study. <i>J Infect Chemother</i> 2018; 24(11): 892-901.	Author
Recent trends in the practice of procedural sedation under local anesthesia for catheter ablation, gastrointestinal endoscopy, and endoscopic surgery in Japan: A retrospective database study in clinical practice from 2012 to 2015. <i>Drugs Real World Outcomes</i> . 2018; 5(3): 137-147.	Author
Low-density lipoprotein cholesterol goal attainment rates by initial statin monotherapy among patients with dyslipidemia and high cardiovascular risk in Japan - A retrospective database analysis. <i>Circ J</i> . 2018; 82(6): 1605-1613.	Acknowledgment
Comorbidities and the use of comedications in people living with HIV on antiretroviral therapy in Japan: a cross-sectional study using a hospital claims database. <i>BMJ Open</i> . 2018; 8(6): e019985.	Acknowledgment
Comorbidities and co-medications in populations with and without chronic hepatitis C virus infection in Japan between 2015 and 2016. <i>BMC Infect Dis</i> . 2018; 18(1): 237.	Acknowledgment
Post-marketing surveillance of donepezil in Japanese patients with Alzheimer's disease using ADAS-J cog: Final reports and constructing a model for ADAS-J cog scores. <i>Japanese Journal of Geriatric Psychiatry</i> . 2018; 29(4): 413-426.	Text (methods)
Pregabalin as an analgesic option for patients undergoing thoracotomy: cost analysis of pregabalin versus epidural analgesia for post-thoracotomy pain relief. <i>J Thorac Dis</i> . 2018; 10(4): 2321-2330.	Author
Development of the Japanese Core Outcome Measures Index (COMI): cross-cultural adaptation and psychometric validation. <i>BMC Musculoskelet Disord</i> . 2018; 19(1): 71.	Author
Long-term safety and efficacy of tofogliflozin as add-on to insulin in patients with type 2 diabetes: Results from a 52-week, multicentre, randomized, double-blind, open-label extension, Phase 4 study in Japan (J-STEP/INS). <i>Diabetes Obes Metab</i> . 2018; 20(5): 1176-1185.	Acknowledgment
Dipeptidyl peptidase-4 inhibitors as preferable oral hypoglycemic agents in terms of treatment satisfaction: Results from a multicenter, 12-week, open label, randomized controlled study in Japan (PREFERENCE 4 study). <i>J Diabetes Investig</i> . 2018; 9(1): 137-145.	Acknowledgment
Decline of rotavirus-coded hospitalizations in children under 5 years: A report from Japan where rotavirus vaccines are self-financed. <i>Vaccine</i> . 2018; 36(19): 2727-2732.	Acknowledgment
Clinical and economic burdens experienced by patients with painful diabetic peripheral neuropathy: An observational study using a Japanese claims database. <i>PLoS One</i> . 2017; 12(10): e0187250.	Author

Assessment of potential risk factors for new onset disabling low back pain in Japanese workers: findings from the CUPID (cultural and psychosocial influences on disability) study. <i>BMC Musculoskelet Disord.</i> 2017; 18(1): 334.	Author
Semi-quantitative Assessment of tumor Spread Through Air Spaces (STAS) in early stage lung adenocarcinomas. <i>J Thorac Oncol.</i> 2017; 12(7): 1046-1051.	Acknowledgment and text (methods)
Development of a Japanese version of the Somatic Symptom Scale-8: Psychometric validity and internal consistency. <i>General Hospital Psychiatry.</i> 2017; 45: 7-11.	Author
Development of the Japanese version of the leeds assessment of the neuropathic symptoms and signs pain scale: Diagnostic utility in a clinical setting. <i>Pain Pract.</i> 2017; 17(6): 800-807.	Author
The Japanese version of the STarT Back Tool predicts 6-month clinical outcomes of low back pain. <i>J Orthop Sci.</i> 2017; 22(2): 224-229.	Author
Usefulness of a simple self-administered joint condition assessment sheet to predict the need for orthopaedic intervention in the management of haemophilic arthropathy. <i>Haemophilia.</i> 2017; 23(2): e116-e123.	Acknowledgment
Psychometric assessment of the Japanese version of the revised fibromyalgia impact questionnaire: reliability and validity. <i>Int J Rheum Dis.</i> 2017; 20(9): 1088-1094.	Author
Comparison of visceral adipose tissue area measured by computed tomography at the different vertebrate level -Sub-analysis using data of Japanese patients derived from International Study of the Prediction of Intra-Abdominal Adiposity and its Relationship with CardioMetabolic Risk/Intra-Abdominal Adiposity (INSPIRE-ME-IAA)-. <i>J Clin Therap Med.</i> 2016; 32(9): 783-791	Acknowledgment
Distribution of ankle-brachial index among inpatients with cardiovascular disease: Analysis using the Kumamoto University Hospital medical database. <i>Annals of vascular diseases,</i> 2016; 9(1): 22-29.	Acknowledgment
Impact of patient attitudes and beliefs to insulin therapy upon initiation, and their attitudinal changes after initiation: the DAWN Japan study. <i>Curr Med Res Opin,</i> 2016; 32(4): 681-686.	Acknowledgment
Host factors and biomarkers associated with poor outcomes in adults with invasive pneumococcal disease. <i>PLoS ONE.</i> 2016; 11(1): e0147877.	Acknowledgment
Pretreatment cognitive profile likely to benefit from donepezil treatment in dementia with lewy bodies: Pooled analyses of two randomized controlled trials. <i>Dement Geriatr Cogn Disord.</i> 2016; 42: 58-68.	Acknowledgment
Development of a Linguistically Validated Japanese Version of the Somatic Symptom Scale-8 (SSS-8). <i>Japanese Journal of Psychosomatic Medicine (Tokyo).</i> 2016; 56(9): 931-937.	Author
Liver-related safety assessment of green tea extracts in humans: a systematic review of randomized controlled trials. <i>EJCN.</i> 2016; 70(11): 1221-1229.	Author
Psychometric Assessment of the Japanese Version of the Zurich Claudication Questionnaire (ZCQ): Reliability and Validity. <i>PLoS ONE.</i> 2016; 11(7): e0160183.	Author
Low back pain prevention exercise – My method for workers –. <i>MB Med Reha.</i> 2016; No.198: 63-69. (in Japanese)	Author
Psychometric Properties of the Japanese Version of the STarT Back Tool in Patients with Low Back Pain. <i>PLoS ONE.</i> 2016; 11(3): e0152019.	Author
Potential risk factors for onset of severe neck and shoulder discomfort (Katakori) in urban Japanese workers. <i>Ind Health.</i> 2016; 54(3): 230-236.	Author
Efficacy and safety assessment of basal supported oral therapy (BOT) with insulin glargine in a real-life clinical setting, stratified by concomitant orally administered antidiabetic agent (OAD) regimens including dipeptidyl peptidase-4 inhibitor (DPP-4i): subanalysis of the ALOHA2 study, drug-use surveillance in Japan. <i>Diabetol Int.</i> 2016; 7(3): 299-307.	Acknowledgment
Predictors for achieving target glycemic control in Japanese patients with type 2 diabetes after initiation of basal supported oral therapy using insulin glargine: sub-analysis of the ALOHA2 study, drug use surveillance in Japan. <i>Diabetol Int.</i> 2016; 7(2): 188-198.	Acknowledgment
Japanese postmarketing surveillance of clopidogrel in patients with non-ST-segment elevation acute coronary syndrome, stable angina, old myocardial infarction, and ST-segment elevation myocardial infarction after percutaneous coronary intervention in a real-life setting: the final report (J-PLACE Final). <i>Cardiovasc Interv Ther.</i> 2016; 31(2): 101-113.	Acknowledgment
The relationship between patient perception of healthcare provision by professionals and the self-care activity of patients with diabetes: Japanese subgroup analysis of the second Diabetes Attitudes, Wishes, and Needs (DAWN2) study. <i>Diabetol Int.</i> 2016; 7(2): 111-118.	Acknowledgment
Disease state changes and safety of long-term donepezil hydrochloride administration in patients with Alzheimer's disease: interim results from the long-term, large-scale J-GOLD study in Japan. <i>Psychogeriatrics.</i> 2016; 2: 107-115.	Author and text (methods)

Development of the Japanese version of the Menorrhagia Multi-Attribute Scale(MMAS): translation and linguistic validation. Sanka to Fujinka. 2015; 82(11): 1299-1305. (in Japanese)	Author
Psychometric properties of the Japanese version of the Tampa Scale for Kinesiophobia (TSK-J) in patients with whiplash neck injury pain and/or low back pain. J Orthop Sci. 2015; 20(6): 985-992.	Author
Can standing back extension exercise improve or prevent low back pain in Japanese care workers? J Man Manip Ther. 2015; 23(4): 205-209.	Author
Adequacy of using consensus guidelines for diagnosis of dementia with lewy bodies in clinical trials for drug development. Dement Geriatr Cogn Disord. 2015; 41(1-2): 55-67.	Acknowledgment
Assessment of psychosocial risk factors for the development of non-specific chronic disabling low back pain in Japanese workers -findings from the Japan Epidemiological Research of Occupation-related Back Pain (JOB) study. Ind Health. 2015; 53(4): 368-377.	Author
Effects of donepezil on extrapyramidal symptoms in patients with dementia with Lewy bodies: a secondary pooled analysis of two randomized-controlled and two open-label long-term extension studies. Dement Geriatr Cogn Disord. 2015; 40(3-4): 186-198.	Acknowledgment
Improved treatment satisfaction and self-reported health status after introduction of basal-supported oral therapy using insulin glargine in patients with type 2 diabetes: sub-analysis of ALOHA2 Study. Diabetes Ther. 2015; 6(2): 153-171.	Acknowledgment
The effects of bazedoxifene on bone, glucose, and lipid metabolism in postmenopausal women with type 2 diabetes: an exploratory pilot study. J Clin Med Res. 2015; 7(10): 762-769.	Acknowledgment
Effect of postoperative administration of pregabalin for post-thoracotomy pain: a randomized study. J Cardiothorac Vasc Anesth. 2015; 29(6): 1567-1572.	Acknowledgment and text (methods)
Epidemiology resesrach on low back pain in Japanese workers. JJOMT. 2015; 63(6): 329-336.	Author
Current situation and issues in titration of basal insulin. Therapeutic Research. 2015; 36(5): 403-411.	Acknowledgment
Linguistic validation of the English version of the Core Lower Urinary Tract Symptom Score. Int J Urol. 2014; 21(12):1291-1292.	Text (methods)
Development of the Japanese version of the Leeds Assessment of Neuropathic Symptoms and Signs Pain Scale: translation and linguistic validation. Pain Clinic. 2014; 35(7): 933-940. (in Japanese)	Author
Psychometric validation of the English version of the overactive bladder symptom score. Urology. 2014; 84(1): 46-50.	Text (methods)
Development of the Japanese version of the Revised Fibromyalgia Impact Questionnaire (JFIQR): translation and linguistic validation. Clinical Rheumatology and Related Research. 2014; 26(1): 35-44. (in Japanese)	Author
Development of the Japanese version of the London Fibromyalgia Epidemiology Study Screening Questionnaire (LFESSQ): translation and linguistic validation.Clinical Rheumatology and Related Research. 2014; 26(2): 130-136. (in Japanese)	Author
Psychometric properties of the Japanese version of the Fear-Avoidance Beliefs Questionnaire (FABQ). J Orthop Sci. 2014; 19(1): 26-32.	Author
Linguistic validation of the English version of the Overactive Bladder Symptom Score. Int J Urol. 2014; 21(2): 229.	Text
Central nervous system effects of the second-generation antihistamines marketed in Japan –review of inter-drug differences using the Proportional Impairment Ratio (PIR)–. PLoS ONE. 2014; 9(12): e114336.	Author
Gaps between actual and desired sex life: web survey of 5,665 Japanese women. J Sex Marital Ther. 2014; 40(1): 33-42.	Text (methods)
Potential risk factors of persistent low back pain developing from mild low back pain in urban Japanese workers. PLoS ONE. 2014; 9(4): e93924.	Author
Validity, reliability and responsiveness of the Japanese version of the Neck Disability Index. J Orthop Sci, 2013; 18(1); 14-21.	Author
Development of a Japanese Version of the Tampa Scale for Kinesiophobia (TSK-J) : Translation and Linguistic Validation. Rinshou Seikeigeka. 2013; 48(1): 13-19. (in Japanese)	Author

Development of the Japanese version of the STarT (Subgrouping for Targeted Treatment) Back Screening Tool; translation and linguistic validation. <i>Journal of Musculoskeletal Pain Research</i> . 2013; 5(1): 11-19. (in Japanese)	Author
Identification of risk factors for new-onset sciatica in Japanese workers: findings from the Japan epidemiological research of Occupation-related Back pain study. <i>Spine (Phila Pa 1976)</i> . 2013; 38(26): E1691-E1700.	Author
Maintenance therapy with pegaptanib sodium for neovascular age-related macular degeneration: an exploratory study in Japanese patients (LEVEL-J study). <i>Jpn J Ophthalmol</i> . 2013; 57(5): 417-423.	Acknowledgment
Development and validation of a new questionnaire assessing quality of life in adults with hypopituitarism: Adult Hypopituitarism Questionnaire (AHQ). <i>PLoS ONE</i> . 2012; 7(9): e44304.	Text (methods)
Development of the Japanese version of the brief symptom inventory-somatization scale: translation and linguistic validation. <i>Seikei Geka</i> . 2012; 63(2): 149-153. (in Japanese)	Author
Potential risk factors for new onset of back pain disability in Japanese workers: findings from the Japan epidemiological research of occupation-related back pain study. <i>Spine (Phila Pa 1976)</i> . 2012; 37(15): 1324-1333.	Author
Recombinant activated factor VII (rFVIIa/NovoSeven®) in the management of severe postpartum haemorrhage: initial report of a multicentre case series in Japan. <i>Int J Hematol</i> . 2012; 95(1): 57-63.	Acknowledgment
Solifenacin as add-on therapy for overactive bladder symptoms in men treated for benign prostatic hyperplasia – ASSIST, randomized controlled study. <i>Urology</i> , 2011; 78(1): 126-33.	Acknowledgment
Development of the Japanese version of the Fear-Avoidance Beliefs Questionnaire (FABQ-J): translation and linguistic validation. <i>Seikei Geka Gakkai</i> . 2011; 62(12): 1301-1306. (in Japanese)	Author
Responsiveness and minimal clinically important change in overactive bladder symptom score. <i>Urology</i> . 2011; 78(4): 768-773.	Acknowledgment
Development of the Japanese version of the Fibromyalgia Impact Questionnaire (JFIQ): psychometric assessments of reliability and validity. <i>Int J Rheum Dis</i> . 2011; 14(1): 74-80.	Author and text (methods)
Comparison of physician's advice for non-specific acute low back pain in Japanese workers: advice to rest versus advice to stay active. <i>Ind Health</i> . 2011; 49(2): 203-208.	Author
Solifenacin as add-on therapy for overactive bladder symptoms in men treated for lower urinary tract symptoms–ASSIST, randomized controlled study. <i>Urology</i> . 2011; 78(1): 126-133.	Acknowledgment
Development of the Japanese version of the Zurich claudication questionnaire (ZCQ): translation and linguistic validation. <i>Nihon Seikei Geka Gakkai</i> . 2010; 61(2): 159-165. (in Japanese)	Author
Development of the Japanese version of the Fibromyalgia Impact Questionnaire (JFIQ): translation and linguistic validation. <i>Clin Rheumatol</i> . 2008; 20: 19-28. (in Japanese)	Author
Opinion survey of women who have male sexual partners on 'better sex'. <i>The Japanese journal of Impotence Research</i> . 2007; 22(3): 287-300. (in Japanese)	Text (methods)
Psychometric properties of the Japanese version of the overactive bladder questionnaire (OAB-q). <i>Journal of the Japan Neurogenic Bladder Society</i> . 2006; 17(2): 250-256. (in Japanese)	Text (methods)

Examples of our experience in supporting conference presentations
(oral or poster)

Title	Conference	Date
Eribulin as first- or second-line chemotherapy for advanced or metastatic HER2-negative breast cancer: a real-world prospective study.	The European Society for Medical Oncology 2018 Congress. Oct 2018, Munich, Germany.	October 19, 2018
The usability of commercial healthcare claims data in Japanese pharmacovigilance: a comparison with a prospective postmarketing study.	The 34th International Conference on Pharmacoepidemiology & Therapeutic Risk Management. Aug 2018. Prague, Czech republic.	August 22, 2018
Treatment patterns and low-density lipoprotein cholesterol goal attainment rates among patients with dyslipidemia and high cardiovascular risk in Japan: a retrospective database analysis	The 82nd Annual Scientific Meeting of the Japanese Circulation Society	March 25, 2018
A real-world observational study of eribulin in patients with soft-tissue sarcoma in Japan.	The 22nd Annual Meeting of Connective Tissue Oncology Society 2017. Nov 2017, Hawaii, US.	November 8, 2017
Prospective observational study of peripheral neuropathy in breast cancer patients treated with eribulin.	The European Society for Medical Oncology 2017 Congress. Sep 2017, Madrid, Spain.	September 11, 2017
Diagnostic capability of "Spine paidetect" to screen for neuropathic pain caused by spinal disorders.	The 44th Annual Meeting of the International Society for the Study of the Lumbar Spine. Jun 2017, Athens, Greece.	June 1, 2017
Effects of donepezil on the prognosis of Parkinson's disease with severe hyposmia - a multicenter randomized double-blind placebo-controlled trial (DASH-PD study).	The 20th International Congress of Parkinson's Disease and Movement Disorders. Jun 2016, Berlin, Germany.	June 22, 2016
Potential risk factors for onset of severe neck and shoulder discomfort (Katakori) in urban Japanese workers.	The 89th Annual Meeting of Japan Society for Occupational Health	May 26, 2016
Psychometric validation of the Japanese version of the Somatic Symptom Scale-8(SSS-8). (in Japanese)	The 8th Annual Meeting of Japanese Association for the Study of Musculoskeletal Pain	December 13, 2015
Association of risk factors for severe low back pain in workers in nursing homes. (in Japanese)	The 23rd Annual Meeting of The Japanese Society of Lumbar Spine Disorders	November 15, 2015
The use of anti-epileptic drugs in Japanese patients with Alzheimer's disease on donepezil: results of large, real-world datasets. (in Japanese)	The 49th Congress of The Japan Epilepsy Society	October 30, 2015
Reality of treatment for patients with Alzheimer's disease in rapidly aging Japanese society: analysis using PMS data of donepezil. (in Japanese)	The 34th Annual Meeting of Japan Society for Dementia Research	October 2, 2015
Analysis of associations between high density lipoprotein structure and cholesterol efflux capacity in dyslipidemic patients treated with pitavastatin (High-Draw study). (in Japanese)	The 47th Annual Meeting of Japan Atherosclerosis Society	July 9, 2015
Gender difference in the progression of Alzheimer's disease in Japanese patients treated with donepezil: Results of the J-GOLD Study.	Alzheimer's Association International Conference 2015. Jul 2015, Washington, US.	July 18, 2015
Risk factors for chronic non-specific low back pain in Japanese workers -JOB Study. (in Japanese)	The 88th Annual Meeting of Japan Society for Occupational Health	May 15, 2015
The potential for liver damage induced by green tea intake: a systematic review of clinical trials.	The 25th Annual Scientific Meeting of the Japanese Epidemiology Association. Jan 2015, Nagoya, Japan.	January 23, 2015
Epidemiological study of low back pain in Japanese workers. (in Japanese)	The 62nd Annual Meeting of Japanese Society of Occupational Medicine and Traumatology symposium 3 "Workers' management of low back pain" (in Japanese)	November 16, 2014
Psychometric properties of the Japanese version of the STarT Back screening tool . (in Japanese)	The 22nd Annual Meeting of The Japanese Society of Lumbar Spine Disorders	November 15, 2014
Potential predictive factors for cognitive response in patients with donepezil treatment: results from the J-GOLD study in Japan.	Alzheimer's Association International Conference 2014. Jul 2014, Copenhagen, Denmark.	July 12, 2014
Impact of early donepezil treatment of Alzheimer's disease in a rapidly aging society: results from the J-GOLD study.	Alzheimer's Association International Conference 2014. Jul 2014, Copenhagen, Denmark.	July 12, 2014

Current outcomes of donepezil treatment in Japan: a comparison between the Phase 3 trial and recent post-marketing (J-GOLD) studies.	Alzheimer's Association International Conference 2014. Jul 2014, Copenhagen, Denmark.	July 12, 2014
Workaholism as a risk factor for depression and disabling back pain among Japanese workers.	The 41st International Society for the Study of the Lumbar Spine. Jun 2014, Seoul, Korea.	June 30, 2014
Usefulness of 'One Stretch', a simple, daily, standing back exercise, for the prevention of onset of aggravation of low back pain in care workers.	The 41st International Society for the Study of the Lumbar Spine. Jun 2014, Seoul, Korea.	June 30, 2014
Identification of risk factors for new-onset sciatica in Japanese Workers: Findings from the Japan Epidemiological Research of Occupation-related Back Pain (JOB) study.	The 23rd International Epidemiology in Occupational Health Conference. Jun 2013, Utrecht, Netherlands.	June 18, 2013
Potential risk factors of persistent low back pain developing from mild LBP in Japanese workers; Findings from a prospective study.	The 40th International Society for the Study of the Lumbar Spine. Apr 2013, Scottsdale, US.	April 30, 2013
Development of the Japanese version of the Revised Fibromyalgia Impact Questionnaire (JFIQR): translation and linguistic validation. (in Japanese)	The 57th Annual General Assembly and Scientific Meeting of the Japan College of Rheumatology	April 18, 2013
Development of the Japanese version of the Tampa Scale for Kinesiophobia (TSK-J): translation and linguistic validation. (in Japanese)	The 5th Annual Meeting of the Japanese Association for the Study of Musculoskeletal Pain	November 17, 2012
Psychometric validation of the Japanese version of the Fear-Avoidance Beliefs Questionnaire (FABQ-J). (in Japanese)	The 20th Annual Meeting of The Japanese Society of Lumbar Spine Disorders	November 13, 2012
Risk factors for new-onset of low back disability among Japanese workers; results from the Cultural and Psychosocial Influences on Disability (CUPID) study.	The 22nd International Epidemiology in Occupational Health Conference. Aug 2011, Oxford, UK.	August 29, 2011
Psychometric validation of the Japanese version of the Neck Disability Index. (in Japanese)	The 40th Annual Meeting of the Japanese Society for Spine Surgery and Related Research	April 22, 2011
Development of the Japanese version of the Fear-Avoidance Beliefs Questionnaire (FABQ): translation and linguistic validation. (in Japanese)	The 18th Annual Meeting of the Japanese Society of Lumbar Spine Disorders	October 31, 2010
Risk factors for new onset of low back disability among Japanese workers; Findings from the Japan epidemiological research of Occupation-related Back pain (JOB) study.	The 21st International Epidemiology in Occupational Health Conference. Apr 2010, Taipei, Taiwan.	April 21, 2010
Development of the Japanese version of the Zurich Claudication Questionnaire (ZCQ): translation and linguistic validation. (in Japanese)	The 17th Annual Meeting of the Japanese Society of Lumbar Spine Disorders	November 21, 2009