

臨床研究および質問票開発の支援実績例 (著者、謝辞、本文中のいずれかに、弊社担当者または弊社名の記載がある公表論文)	
標題・掲載雑誌	記載箇所
Characteristics of patients with longer treatment period of lenvatinib for unresectable hepatocellular carcinoma: a post-hoc analysis of post-marketing surveillance study in Japan. PLOS ONE. 2024; 19(3): e0298420.	謝辞
Association between glycemc control and patient-reported outcomes in adults with type 1 diabetes in Japan: the SAGE study subanalysis. Diabetol Int. 2023; 15(2): 212-222.	謝辞
Validation and comparison of the coding algorithms to identify people with migraine using Japanese claims data. Front Neurol. 2023; 14: 1231351.	謝辞
Real-world treatment patterns of subsequent therapy after palbociclib in patients with advanced breast cancer in Japan. Breast. 2023; 70: 1-7.	謝辞
Real-world treatment patterns of novel drugs in relapsed or refractory acute lymphoblastic leukemia patients in Japan. Future Oncol. 2023; 19(19): 1343-1356.	謝辞
Exploring changes in attitudes, behaviors, and self-measured health data through lifestyle modification support by community pharmacists: Suito-Ogaki Selfcare (SOS) trial. Integr Pharm Res Pract. 2023; 12: 87-99.	謝辞
Impact of tolvaptan add-on treatment on patients with heart failure requiring long-term congestion management: a retrospective cohort study using a medical claim database in Japan. J Cardiol. 2023; 82(1): 35-42.	謝辞
Perioperative analgesics and anesthesia as risk factors for postoperative chronic opioid use in patients undergoing total knee arthroplasty: a retrospective cohort study using Japanese hospital claims data. Drugs - Real World Outcomes. 2023; 10(2): 331-340.	著者、謝辞
Safety and effectiveness of molnupiravir (LAGEVRIO®) capsules in Japanese patients with COVID-19: interim report of post-marketing surveillance in Japan. Infect Dis Ther. 2023; 12(4): 1119-1136.	謝辞
Validation of algorithms to identify colorectal cancer patients from administrative claims data of a Japanese hospital. BMC Health Serv Res. 2023; 23(1): 274.	著者、謝辞
Linguistic and psychometric validation of the cognition bolt-on version of the Japanese EQ-5D-5L for the elderly. J Alzheimers Dis. 2023; 91(4): 1447-1458.	著者、謝辞
Clinical characteristics, medication use, and impact of primary headache on daily activities: an observational study using linked online survey and medical claims data in Japan. BMC Neurol. 2023; 23(1): 80.	謝辞
An attempt to replicate randomized trials of diabetes treatments using a Japanese administrative claims and health checkup database: a feasibility study. Drugs - Real World Outcomes. 2023; 10(2): 235-247.	著者、謝辞
Validation of algorithms to identify bone metastases using administrative claims data in a Japanese hospital. Drugs - Real World Outcomes. 2023; 10(2) 187-194.	著者、謝辞
Safety and effectiveness of lenvatinib in patients with unresectable hepatocellular carcinoma in real-world clinical practice: an observational post-marketing study in Japan. Drugs - Real World Outcomes. 2023; 10(2): 195-205.	謝辞

Impact of "time zero" of follow-up settings in a comparative effectiveness study using real-world data with a non-user comparator: comparison of six different settings. <i>Drugs - Real World Outcomes</i> . 2023; 10(1): 107-117.	著者、謝辞
Cardiometabolic risk reductions in patients with type 2 diabetes mellitus newly treated with a sodium–glucose cotransporter 2 inhibitor versus a dipeptidyl peptidase-4 inhibitor: A real-world administrative database study in Japan. <i>J Diabetes Investig</i> . 2023; 14(3): 404-416.	著者、謝辞
Physician consultation rates and characteristics among workers with chronic pain or headache who participated in a behavioural change program: a retrospective database analysis using real-world healthcare data. <i>BMJ Open</i> . 2022; 12(11): e056846.	謝辞
Effect of parenteral energy or amino acid doses on in-hospital mortality among patients with aspiration pneumonia: a cohort medical claims database study. <i>J Gerontol A Biol Sci Med Sci</i> . 2022; 77(8): 1683-1690.	謝辞
Treatment discontinuation among patients with schizophrenia treated with brexpiprazole and other oral atypical antipsychotics in Japan: A retrospective observational study. <i>Adv Ther</i> . 2022; 39(9): 4299-4314.	謝辞
Reduction in asthma exacerbation rate after mepolizumab treatment initiation in patients with severe asthma: A real-world database study in Japan. <i>Pulm Pharmacol Ther</i> . 2022; 75: 102130.	謝辞
Effect of donepezil for dementia prevention in Parkinson's disease with severe hyposmia (The DASH-PD study): A randomized long-term placebo-controlled trial. <i>eClinicalMedicine</i> . 2022; 51: 101571.	著者
The use of a network analysis to identify associations and temporal patterns among non-communicable diseases in Japan based on a large medical claims database. <i>Drugs - Real World Outcomes</i> . 2022; 9(3): 463-476.	著者、資金提供
A study to investigate the prevalence of headache disorders and migraine among people registered in a health insurance association in Japan. <i>J Headache Pain</i> . 2022; 23(1): 70.	謝辞
Association between HbA1c levels and diabetic peripheral neuropathy: A case–control study of patients with type 2 diabetes using claims data. <i>Drugs - Real World Outcomes</i> . 2022; 9(3): 403-414.	謝辞
C型慢性肝炎・代償性肝硬変患者におけるグレカプレビル水和物/ピブレンタスビル配合剤の安全性と有効性：使用成績調査に基づいた検討. <i>肝臓</i> . 2022; 63(3): 120-150.	謝辞
Real-world treatment patterns of palbociclib and blood count monitoring in patients with advanced breast cancer in Japan. <i>Future Oncol</i> . 2022; 18(17): 2101-2111.	謝辞
Comparative analysis of medical costs after hepatectomy versus radiofrequency ablation in patients with hepatocellular carcinoma in real-world clinical practice. <i>Hepatol Res</i> . 2022; 52(5): 471-478.	謝辞
腰痛と坐骨神経痛に対する薬物療法に関するナラティブレビュー. <i>J Spine Res</i> . 2022; 13(2): 67-86.	謝辞
健常成人を対象としたEDC, ePRO, およびウェアラブルデバイスを用いた共同調査結果とその有用性. <i>Prog Med</i> . 2022; 42(1): 91-100.	謝辞
Internet survey on factors associated with care-seeking behaviours of people with chronic musculoskeletal pain in Japan. <i>J Orthop Surg (Hong Kong)</i> . 2021; 29(3): 23094990211044836.	資金提供
A post marketing survey of intravenous daptomycin in Japanese patients. <i>J. New Rem. & Clin</i> . 2021; 70: 1189-1215.	謝辞

Nutritional management in inpatients with aspiration pneumonia: a cohort medical claims database study. Arch Gerontol Geriatr. 2021; 95: 104398.	謝辞
Treatment patterns and medical costs after hepatectomy in real-world practice for patients with hepatocellular carcinoma in Japan. Hepatol Res. 2021; 51(10): 1073-1081.	謝辞
Understanding confidence in the human papillomavirus vaccine in Japan: a web-based survey of mothers, female adolescents, and healthcare professionals. Hum Vaccin Immunother. 2021; 17(9): 3102-3112.	謝辞
Real-world treatment patterns of patients with atopic dermatitis in Japan: analysis of the JMDC claims database. J Cutan Immunol Allergy. 2021; 00: 1-11.	謝辞
Multiple comorbidities increase the risk of death from invasive pneumococcal disease under the age of 65 years. J Infect Chemother. 2021; 27(9): 1311-1318.	謝辞
Glycemic control status, diabetes management patterns, and clinical characteristics of adults with type 1 diabetes in Japan: study of adults' glycemia in T1DM subanalysis. Diabetol Int. 2021; 12(4): 460-473.	謝辞
Real-world treatment patterns and adherence to oral medication among patients with bipolar disorders: a retrospective, observational study using a healthcare claims database. Neuropsychiatr Dis Treat. 2021; 17: 821-833.	謝辞
Real-world hospital mortality of liver cirrhosis inpatients in Japan: a large-scale cohort study using a medical claims database: prognosis of liver cirrhosis. Hepatol Res. 2021; 51(6): 682-693.	謝辞
Effectiveness of pregabalin treatment for neuropathic pain in patients with spine diseases: a pooled analysis of two multicenter observational studies in Japan. J Pain Res. 2021; 14: 757-771.	謝辞
Antidyslipidemic drug prescriptions and lipid control status after unfavorable annual health checkup results: a retrospective cohort study using a health insurance database. Drugs - Real World Outcomes. 2021; 8(2): 227-240.	謝辞
Indices of peripheral leukocytes predict longer overall survival in breast cancer patients on eribulin in Japan. Breast Cancer. 2021; 28(4): 945-955.	謝辞
Changes in the penetration rate of biosimilar infliximab within Japan using a Japanese claims database. Clinicoecon Outcomes Res. 2021; 13: 145-153.	著者
Relationship of annual health checkups with the subject's subsequent behavior of cardiovascular risk management in a real-world setting in Japan: a retrospective cohort study on changes in antihypertensive drug prescription and blood pressure from 2015 to 2017. Drugs Real World Outcomes. 2021; 8(2): 215-225.	謝辞
Risk of hypoglycemia in Japanese people with type 2 diabetes mellitus who initiated or switched to insulin glargine 300 U/mL: a subgroup analysis of 12-month post-marketing surveillance study (X-STAR study). Diabetes Res Clin Pract. 2021; 172: 108647.	謝辞
Clinical impact of prescribed doses of nutrients for patients exclusively receiving parenteral nutrition in Japanese hospitals: a retrospective cohort study. J Parenter Enteral Nutr. 2021; 45(7): 1514-1522.	謝辞
Treatment and healthcare cost among patients with hip or knee osteoarthritis: a cross-sectional study using a real-world claims database in Japan between 2013 and 2019. Clin Drug Investig. 2020; 40(11): 1071-1084.	著者, 謝辞
Comparison of hospital length of stay of acute ischemic stroke patients with non-valvular atrial fibrillation started on rivaroxaban or warfarin treatment during hospitalization. J Med Econ. 2020; 23(12): 1379-1388.	著者, 謝辞

Real-world data on the use of insulin glargine 300 U/mL in Japanese patients with type 1 diabetes: twelve-month results from a post-marketing surveillance study (X-STAR study). <i>Expert Opin Pharmacother.</i> 2021; 22(2): 249-256.	謝辞
Effectiveness and safety of insulin glargine 300 unit/mL in Japanese type 2 diabetes mellitus patients: a 12-month post-marketing surveillance study (X-STAR study). <i>Expert Opin Pharmacother.</i> 2020; 21: 1771-1780.	謝辞
A survey on total parenteral nutrition in 55,000 hospitalized patients: retrospective cohort study using a medical claims database. <i>Clin Nutr ESPEN.</i> 2020; 39:198-205.	著者, 謝辞
Safety and effectiveness of lenvatinib in 594 patients with unresectable thyroid cancer in an all-case post-marketing observational study in Japan. <i>Adv Ther.</i> 2020; 37(9): 3850-3862.	著者
Psychometric properties of the Japanese version of the Kansas City Cardiomyopathy Questionnaire in Japanese patients with chronic heart failure. <i>Health Qual Life Outcomes.</i> 2020; 18(1): 236.	著者
Incidence of and risk factors for recurrent <i>Clostridioides difficile</i> infection in Japan using a claims database: a retrospective cohort study. <i>Anaerobe.</i> 2020; 61: 102139.	謝辞
肝硬変入院患者の腹水合併および治療の実態調査：診療データベースを用いたコホート研究. <i>肝臓.</i> 2020; 61(6): 314-325.	謝辞
Treatment persistence between long-acting injectable versus orally administered aripiprazole among patients with schizophrenia in a real-world clinical setting in Japan. <i>Adv Ther.</i> 2020; 37(7): 3324-3336.	謝辞
Effectiveness and safety of eribulin in Japanese patients with HER2-negative, advanced breast cancer: a 2-year post-marketing observational study in a real-world setting. <i>Invest New Drugs.</i> 2020; 38(5): 1540-1549.	謝辞
Long-term tolerability and effectiveness of raltegravir in Japanese patients: results from post-marketing surveillance. <i>PLoS One.</i> 2019; 14(1): e0210384.	謝辞
Patterns of drug treatment in patients with osteoarthritis and chronic low back pain in Japan: a retrospective database study. <i>J Pain Res.</i> 2019; 12: 1631-1648.	謝辞
Daily practice and prognostic factors for pneumonia caused by methicillin-resistant <i>Staphylococcus aureus</i> in Japan: a multicenter prospective observational cohort study. <i>J Infect Chemother.</i> 2020; 26(2): 242-251.	謝辞
皮膚外用剤の物性の違いが塗布量に与える影響に関する臨床研究：成人被験者を対象としたアンケート調査による基剤の使用感と塗布量の探索的検討. <i>Yakugaku Zasshi.</i> 2019; 139(10): 1313-1325.	謝辞
Development of a Japanese version of the Örebro Musculoskeletal Pain Screening Questionnaire (ÖMPSQ-J) and the short form version: translation and linguistic validation. <i>Journal of Musculoskeletal Pain Research.</i> 2019; 11: 35-48.	著者
A cost-effectiveness analysis of pregabalin for the treatment of patients with chronic cervical pain with a neuropathic component in Japan. <i>J Pain Res.</i> 2019; 12: 2785-2797.	著者
Safety and effectiveness of tofogliflozin in elderly Japanese patients with type 2 diabetes mellitus: a subanalysis of a post-marketing study (J-STEP/EL Study). <i>J Diabetes Investig.</i> 2020; 11(2): 405-416.	謝辞
Sustained reduction in rotavirus-coded hospitalizations in children aged <5 years after introduction of self-financed rotavirus vaccines in Japan. <i>Hum Vaccin Immunother.</i> 2020; 16(1): 132-137.	謝辞

Safety and effectiveness of tofogliflozin in Japanese patients with type 2 diabetes mellitus in real-world practice: results of 12-month interim analysis of a long-term post-marketing surveillance study (J-STEP/LT). <i>J Diabetes Investig.</i> 2020; 11(1): 132-141.	謝辞
Switch rates, time-to-switch, and switch patterns of antiretroviral therapy in people living with human immunodeficiency virus in Japan, in a hospital-claim database. <i>BMC Infect Dis.</i> 2019; 19(1): 505.	謝辞
Pharmacotherapy decision-making among patients with breast cancer in Japan: results of an online survey. <i>Breast Cancer.</i> 2019; 26(6): 799-807.	謝辞
特定健康診査の受診率に関連する社会人口学的因子および施策の検討. <i>厚生の指標.</i> 2019; 66(4): 29-35.	謝辞
Use of a healthcare claims database for post-marketing safety assessments of eribulin in Japan: a comparative assessment with a prospective post-marketing surveillance study. <i>Drugs Real World Outcomes.</i> 2019; 6(1): 27-35.	著者
Secular trend in the incidence of Japanese employees with thyroid cancer undergoing thyroidectomy from 2005 to 2014: a retrospective descriptive study using an employment-based insurance claims database. <i>J of thyroid cancer.</i> 2019; 1: 5-15.	-
Ultra-high-dose methylcobalamin in amyotrophic lateral sclerosis: a long-term phase II/III randomised controlled study. <i>J Neurol Neurosurg Psychiatry.</i> 2019; 90(4): 451-457.	謝辞
Differences in the incidence of postoperative pneumonia after percutaneous endoscopic gastrostomy between liquid and semi-solid nutrient administration. <i>Eur J Clin Nutr.</i> 2019; 73(2): 250-257.	謝辞, 本文 (方法)
Assessment of risk factors for non-specific chronic disabling low back pain in Japanese workers—findings from the CUPID (Cultural and Psychosocial Influences on Disability) study. <i>Industrial Health.</i> 2019; 57(4): 503-510.	著者
Greater burden of chronic comorbidities and co-medications among people living with HIV versus people without HIV in Japan: a hospital claims database study. <i>J Infect Chemother.</i> 2019; 25(2): 89-95.	謝辞
Chemotherapy-induced peripheral neuropathy in breast cancer patients treated with eribulin: interim data from a post-marketing observational study. <i>Breast Cancer.</i> 2019; 26(2): 235-243.	謝辞
心因性疼痛を考える：用語としての認知性疼痛の提案. <i>Pain Research.</i> 2018; 33(3): 183-192.	利益相反
Healthcare burden of recurrent <i>Clostridioides difficile</i> infection in Japan: a retrospective database study. <i>J Infect Chemother.</i> 2018; 24(11): 892-901.	著者
Recent trends in the practice of procedural sedation under local anesthesia for catheter ablation, gastrointestinal endoscopy, and endoscopic surgery in Japan: a retrospective database study in clinical practice from 2012 to 2015. <i>Drugs Real World Outcomes.</i> 2018; 5(3): 137-147.	著者
Low-density lipoprotein cholesterol goal attainment rates by initial statin monotherapy among patients with dyslipidemia and high cardiovascular risk in Japan—a retrospective database analysis. <i>Circ J.</i> 2018; 82(6): 1605-1613.	謝辞
Comorbidities and the use of comedications in people living with HIV on antiretroviral therapy in Japan: a cross-sectional study using a hospital claims database. <i>BMJ Open.</i> 2018; 8(6): e019985.	謝辞
Comorbidities and co-medications in populations with and without chronic hepatitis C virus infection in Japan between 2015 and 2016. <i>BMC Infect Dis.</i> 2018; 18(1): 237.	謝辞

アルツハイマー型認知症に対するドネペジル塩酸塩によるADAS-J cogを用いた調査—アリセプト®特定使用成績調査最終報告とADAS-J cog合計得点推移モデルの構築—. 老年精神医学雑誌. 2018; 29(4): 413-426.	本文 (方法)
Pregabalin as an analgesic option for patients undergoing thoracotomy: cost analysis of pregabalin versus epidural analgesia for post-thoracotomy pain relief. J Thorac Dis. 2018; 10(4): 2321-2330.	著者
Development of the Japanese Core Outcome Measures Index (COMI): cross-cultural adaptation and psychometric validation. BMC Musculoskelet Disord. 2018; 19(1): 71.	著者
Long-term safety and efficacy of tofogliflozin as add-on to insulin in patients with type 2 diabetes: results from a 52-week, multicentre, randomized, double-blind, open-label extension, Phase 4 study in Japan (J-STEP/INS). Diabetes Obes Metab. 2018; 20(5): 1176-1185.	謝辞
Dipeptidyl peptidase-4 inhibitors as preferable oral hypoglycemic agents in terms of treatment satisfaction: results from a multicenter, 12-week, open label, randomized controlled study in Japan (PREFERENCE 4 study). J Diabetes Investig. 2018; 9(1): 137-145.	謝辞
Decline of rotavirus-coded hospitalizations in children under 5 years: a report from Japan where rotavirus vaccines are self-financed. Vaccine. 2018; 36(19): 2727-2732.	謝辞
Clinical and economic burdens experienced by patients with painful diabetic peripheral neuropathy: an observational study using a Japanese claims database. PLoS One. 2017; 12(10): e0187250.	著者
Assessment of potential risk factors for new onset disabling low back pain in Japanese workers: findings from the CUPID (cultural and psychosocial influences on disability) study. BMC Musculoskelet Disord. 2017; 18(1): 334.	著者
Semi-quantitative assessment of tumor spread through air spaces (STAS) in early stage lung adenocarcinomas. J Thorac Oncol. 2017; 12(7): 1046-1051.	謝辞, 本文 (方法)
Development of a Japanese version of the Somatic Symptom Scale-8: psychometric validity and internal consistency. General Hospital Psychiatry. 2017; 45: 7-11.	著者
Development of the Japanese version of the leeds assessment of the neuropathic symptoms and signs pain scale: diagnostic utility in a clinical setting. Pain Pract. 2017; 17(6): 800-807.	著者
The Japanese version of the STarT Back Tool predicts 6-month clinical outcomes of low back pain. J Orthop Sci. 2017; 22(2): 224-229.	著者
Usefulness of a simple self-administered joint condition assessment sheet to predict the need for orthopaedic intervention in the management of haemophilic arthropathy. Haemophilia. 2017; 23(2): e116-e123.	謝辞
Psychometric assessment of the Japanese version of the revised fibromyalgia impact questionnaire: reliability and validity. Int J Rheum Dis. 2017; 20(9): 1088-1094.	著者
腹部コンピューター断層撮影による内臓脂肪面積測定の国際比較—INternational Study of the Prediction of Intra-Abdominal Adiposity and its RELationship with CardioMEtabolic Risk/Intra-Abdominal Adiposity (INSPIRE-ME-IAA) の日本人のデータを使用した横断研究—. 臨床医薬. 2016; 32(9): 783-791.	謝辞
Distribution of ankle-brachial index among inpatients with cardiovascular disease: analysis using the Kumamoto University Hospital medical database. Annals of vascular diseases, 2016; 9(1): 22-29.	謝辞
Impact of patient attitudes and beliefs to insulin therapy upon initiation, and their attitudinal changes after initiation: the DAWN Japan study. Curr Med Res Opin. 2016; 32(4): 681-686.	謝辞

Host factors and biomarkers associated with poor outcomes in adults with invasive pneumococcal disease. PLoS ONE. 2016; 11(1): e0147877.	謝辞
Pretreatment cognitive profile likely to benefit from donepezil treatment in dementia with lewy bodies: pooled analyses of two randomized controlled trials. Dement Geriatr Cogn Disord. 2016; 42: 58-68.	謝辞
日本語版Somatic Symptom Scale-8 (SSS-8 [身体症状スケール]) の開発—言語的妥当性を担保した翻訳版の作成—. 心身医学. 2016; 56(9): 931-937.	著者
Liver-related safety assessment of green tea extracts in humans: a systematic review of randomized controlled trials. EJCN. 2016; 70(11): 1221-1229.	著者
Psychometric assessment of the Japanese version of the Zurich Claudication Questionnaire (ZCQ): reliability and validity. PLoS ONE. 2016; 11(7): e0160183.	著者
腰痛予防のエクササイズ—労働者に対する私の方法—. MB Med Reha. 2016; No.198: 63-69.	著者
Psychometric properties of the Japanese version of the STarT Back tool in patients with low back pain. PLoS ONE. 2016; 11(3): e0152019.	著者
Potential risk factors for onset of severe neck and shoulder discomfort (Katakori) in urban Japanese workers. Ind Health. 2016; 54(3): 230-236.	著者
Efficacy and safety assessment of basal supported oral therapy (BOT) with insulin glargine in a real-life clinical setting, stratified by concomitant orally administered antidiabetic agent (OAD) regimens including dipeptidyl peptidase-4 inhibitor (DPP-4i): subanalysis of the ALOHA2 study, drug-use surveillance in Japan. Diabetol Int. 2016; 7(3): 299-307.	謝辞
Predictors for achieving target glycemic control in Japanese patients with type 2 diabetes after initiation of basal supported oral therapy using insulin glargine: sub-analysis of the ALOHA2 study, drug use surveillance in Japan. Diabetol Int. 2016; 7(2): 188-198.	謝辞
Japanese postmarketing surveillance of clopidogrel in patients with non-ST-segment elevation acute coronary syndrome, stable angina, old myocardial infarction, and ST-segment elevation myocardial infarction after percutaneous coronary intervention in a real-life setting: the final report (J-PLACE Final). Cardiovasc Interv Ther. 2016; 31(2): 101-113.	謝辞
The relationship between patient perception of healthcare provision by professionals and the self-care activity of patients with diabetes: Japanese subgroup analysis of the second Diabetes Attitudes, Wishes, and Needs (DAWN2) study. Diabetol Int. 2016; 7(2): 111-118.	謝辞
Disease state changes and safety of long-term donepezil hydrochloride administration in patients with Alzheimer's disease: interim results from the long-term, large-scale J-GOLD study in Japan. Psychogeriatrics. 2016; 2: 107-115.	著者, 本文 (方法)
日本語版Menorrhagia Multi-Attribute Scale (MMAS) の開発 : 言語的妥当性の検討. 産科と婦人科. 2015; 82(11): 1299-1305.	著者
Psychometric properties of the Japanese version of the Tampa Scale for Kinesiophobia (TSK-J) in patients with whiplash neck injury pain and/or low back pain. J Orthop Sci. 2015; 20(6): 985-992.	著者
Can standing back extension exercise improve or prevent low back pain in Japanese care workers? J Man Manip Ther. 2015; 23(4): 205-209.	著者
Adequacy of using consensus guidelines for diagnosis of dementia with lewy bodies in clinical trials for drug development. Dement Geriatr Cogn Disord. 2015; 41(1-2): 55-67.	謝辞

Assessment of psychosocial risk factors for the development of non-specific chronic disabling low back pain in Japanese workers—findings from the Japan Epidemiological Research of Occupation-related Back Pain (JOB) study. <i>Ind Health</i> . 2015; 53(4): 368-377.	著者
Effects of donepezil on extrapyramidal symptoms in patients with dementia with Lewy bodies: a secondary pooled analysis of two randomized-controlled and two open-label long-term extension studies. <i>Dement Geriatr Cogn Disord</i> . 2015; 40(3-4): 186-198.	謝辞
Improved treatment satisfaction and self-reported health status after introduction of basal-supported oral therapy using insulin glargine in patients with type 2 diabetes: sub-analysis of ALOHA2 Study. <i>Diabetes Ther</i> . 2015; 6(2): 153-171.	謝辞
The effects of bazedoxifene on bone, glucose, and lipid metabolism in postmenopausal women with type 2 diabetes: an exploratory pilot study. <i>J Clin Med Res</i> . 2015; 7(10): 762-769.	謝辞
Effect of postoperative administration of pregabalin for post-thoracotomy pain: a randomized study. <i>J Cardiothorac Vasc Anesth</i> . 2015; 29(6): 1567-1572.	謝辞, 本文 (方法)
日本人勤労者を対象とした腰痛疫学研究. <i>日本職業・災害医学会会誌</i> . 2015; 63(6): 329-336.	著者
基礎インスリンの用量調節に関する現状と課題. <i>Therapeutic Research</i> . 2015; 36(5): 403-411.	謝辞
Linguistic validation of the English version of the Core Lower Urinary Tract Symptom Score. <i>Int J Urol</i> . 2014; 21(12): 1291-1292.	本文 (方法)
日本語版Leeds Assessment of Neuropathic Symptoms and Signs Pain Scale (LANSS痛みスケール) の開発: 言語的妥当性を担保した翻訳版の作成. <i>ペインクリニック</i> . 2014; 35(7): 933-940.	著者
Psychometric validation of the English version of the overactive bladder symptom score. <i>Urology</i> . 2014; 84(1): 46-50.	本文 (方法)
Revised Fibromyalgia Impact Questionnaire (FIQR) 日本語版の開発: 言語的妥当性を担保した翻訳版の作成. <i>臨床リウマチ</i> . 2014; 26(1): 35-44.	著者
日本語版London Fibromyalgia Epidemiology Study Screening Questionnaire (LFESSQ) の開発: 言語的妥当性を担保した翻訳版の作成. <i>臨床リウマチ</i> . 2014; 26(2): 130-136.	著者
Psychometric properties of the Japanese version of the Fear-Avoidance Beliefs Questionnaire (FABQ). <i>J Orthop Sci</i> . 2014; 19(1): 26-32.	著者
Linguistic validation of the English version of the Overactive Bladder Symptom Score. <i>Int J Urol</i> . 2014; 21(2): 229.	本文
Central nervous system effects of the second-generation antihistamines marketed in Japan—review of inter-drug differences using the Proportional Impairment Ratio (PIR)—. <i>PLoS ONE</i> . 2014; 9(12): e114336.	著者
Gaps between actual and desired sex life: web survey of 5,665 Japanese women. <i>J Sex Marital Ther</i> . 2014; 40(1): 33-42.	本文 (方法)
Potential risk factors of persistent low back pain developing from mild low back pain in urban Japanese workers. <i>PLoS ONE</i> . 2014; 9(4): e93924.	著者

Validity, reliability and responsiveness of the Japanese version of the Neck Disability Index. J Orthop Sci, 2013; 18(1); 14-21.	著者
日本語版Tampa Scale for Kinesiophobia (TSK-J) の開発：言語的妥当性を担保した翻訳版の作成. 臨床整形外科. 2013; 48(1): 13-19.	著者
日本語版STarT (Subgrouping for Targeted Treatment) Backスクリーニングツールの開発—言語的妥当性を担保した翻訳版の作成—. 日本運動器疼痛学会誌. 2013; 5(1): 11-19.	著者
Identification of risk factors for new-onset sciatica in Japanese workers: findings from the Japan epidemiological research of Occupation-related Back pain study. Spine (Phila Pa 1976). 2013; 38(26): E1691-E1700.	著者
Maintenance therapy with pegaptanib sodium for neovascular age-related macular degeneration: an exploratory study in Japanese patients (LEVEL-J study). Jpn J Ophthalmol. 2013; 57(5): 417-423.	謝辞
Development and validation of a new questionnaire assessing quality of life in adults with hypopituitarism: Adult Hypopituitarism Questionnaire (AHQ). PLoS ONE. 2012; 7(9): e44304.	本文 (方法)
創意と工夫 日本語版brief symptom inventory-somatizationの開発：言語的妥当性を担保した翻訳版の作成. 整形外科. 2012; 63(2): 149-153.	著者
Potential risk factors for new onset of back pain disability in Japanese workers: findings from the Japan epidemiological research of occupation-related back pain study. Spine (Phila Pa 1976). 2012; 37(15): 1324-1333.	著者
Recombinant activated factor VII (rFVIIa/NovoSeven®) in the management of severe postpartum haemorrhage: initial report of a multicentre case series in Japan. Int J Hematol. 2012; 95(1): 57-63.	謝辞
日本語版Fear-Avoidance Beliefs Questionnaire (FABQ-J) の開発—言語的妥当性を担保した翻訳版の作成—. 整形外科. 2011; 62(12): 1301-1306.	著者
Responsiveness and minimal clinically important change in overactive bladder symptom score. Urology. 2011; 78(4): 768-773.	謝辞
Development of the Japanese version of the Fibromyalgia Impact Questionnaire (JFIQ): psychometric assessments of reliability and validity. Int J Rheum Dis. 2011; 14(1): 74-80.	著者, 本文 (方法)
Comparison of physician's advice for non-specific acute low back pain in Japanese workers: advice to rest versus advice to stay active. Ind Health. 2011; 49(2): 203-208.	著者
Solifenacin as add-on therapy for overactive bladder symptoms in men treated for lower urinary tract symptoms—ASSIST, randomized controlled study. Urology. 2011; 78(1): 126-133.	謝辞
日本語版Zurich claudication questionnaire (ZCQ) の開発—言語的妥当性を担保した翻訳版の作成—. 整形外科. 2010; 61(2): 159-165.	著者
日本語版Fibromyalgia Impact Questionnaire (JFIQ) の開発：言語的妥当性を担保した翻訳版の作成. 臨床リウマチ. 2008; 20: 19-28.	著者
女性を対象とした「よりよい性生活」に関する意識調査. 日本性機能学会雑誌. 2007; 22(3): 287-300.	本文 (方法)
Overactive bladder questionnaire (OAB-q) の日本語版の計量心理学的検討. 日本排尿機能学会誌. 2006; 17(2): 250-256.	本文 (方法)

学会発表の支援実績例
(口頭およびポスター発表)

標題	学会名	発表年月日
Impact of primary headache on daily activities from real world data in Japan	IHC 2023 - International Headache Congress; Seoul, Korea	2023年9月14日
Linguistic and psychometric validation of the "cognition" bolt-on version of the Japanese EQ-5D-5L (EQ-5D-5L+C) by adding a cognitive dimension for the elderly	ISPOR Europe 2022; Vienna, Austria	2022年11月7日
Safety and efficacy of long-term treatment with perampanel in Japanese patients with partial-onset seizure or primary generalized tonic-clonic seizure in a real-world setting	American Epilepsy Society Annual Meeting 2019; Baltimore, US	2019年12月9日
変形性関節症および慢性腰痛患者の薬物治療実態と変形性関節症に関連した手術を受けた患者の特徴	第12回日本運動器疼痛学会	2019年11月30日
インスリン未治療 2 型糖尿病患者におけるインスリングルルギン 300U/mL 入院導入の有効性と安全性の検討: X-STAR studyサブ解析	第62回日本糖尿病学会年次学術集会	2019年5月23日
Eribulin as first- or second-line chemotherapy for advanced or metastatic HER2-negative breast cancer: a real-world prospective study	The European Society for Medical Oncology 2018 Congress; Munich, Germany	2018年10月19日
The usability of commercial healthcare claims data in Japanese pharmacovigilance: a comparison with a prospective postmarketing study	The 34th International Conference on Pharmacoepidemiology & Therapeutic Risk Management; Prague, Czech republic	2018年8月22日
Treatment patterns and low-density lipoprotein cholesterol goal attainment rates among patients with dyslipidemia and high cardiovascular risk in Japan: a retrospective database analysis	第82回日本循環器学会学術集会	2018年3月25日
A real-world observational study of eribulin in patients with soft-tissue sarcoma in Japan	The 22nd Annual Meeting of Connective Tissue Oncology Society 2017; Hawaii, US	2017年11月8日
Prospective observational study of peripheral neuropathy in breast cancer patients treated with eribulin	The European Society for Medical Oncology 2017 Congress; Madrid, Spain	2017年9月11日
Diagnostic capability of "Spine paidetect" to screen for neuropathic pain caused by spinal disorders	The 44th Annual Meeting of the International Society for the Study of the Lumbar Spine; Athens, Greece	2017年6月1日
Effects of donepezil on the prognosis of Parkinson's disease with severe hyposmia—a multicenter randomized double-blind placebo-controlled trial (DASH-PD study)	The 20th International Congress of Parkinson's Disease and Movement Disorders; Berlin, Germany	2016年6月22日
日本人勤労者における重度の肩こり発生に関連する危険因子の検討	第89回日本産業衛生学会	2016年5月26日
日本語版Somatic Symptom Scale-8 (SSS-8) の計量心理学的な信頼性と妥当性の検討	第8回日本運動器疼痛学会	2015年12月13日
介護施設職員における重症度の高い腰痛の関連要因の検討	第23回日本腰痛学会	2015年11月15日

日本のアルツハイマー型認知症患者における抗てんかん薬の使用実態：ドネペジルのPMSデータを用いた解析	第49回日本てんかん学会学術集会	2015年10月30日
超高齢化社会日本でのAD患者の治療実態：ドネペジルのPMSデータを用いた解析	第34回日本認知症学会学術集会	2015年10月2日
スタチン投与によるコレステロール引き抜き能とHDL亜分画ならびにHDL構成タンパクへの影響	第47回日本動脈硬化学会	2015年7月9日
Gender difference in the progression of Alzheimer's disease in Japanese patients treated with donepezil: Results of the J-GOLD Study	Alzheimer's Association International Conference 2015; Washington, US	2015年7月18日
日本人勤労者における慢性非特異的腰痛の危険因子勤労者：JOB Studyを用いた検討	第88回日本産業衛生学会	2015年5月15日
緑茶摂取量によって誘発される肝障害の可能性:臨床試験の系統的レビュー	第25回日本疫学会学術総会	2015年1月23日
日本人勤労者を対象とした腰痛疫学研究	第62回日本職業・災害医学会学術大会, シンポジウム3「勤労者の腰痛管理」	2014年11月16日
日本語版 STarT Back スクリーニングツールの計量心理学的検討	第22回日本腰痛学会	2014年11月15日
Potential predictive factors for cognitive response in patients with donepezil treatment: results from the J-GOLD study in Japan	Alzheimer's Association International Conference 2014; Copenhagen, Denmark	2014年7月12日
Impact of early donepezil treatment of Alzheimer's disease in a rapidly aging society: results from the J-GOLD study	Alzheimer's Association International Conference 2014; Copenhagen, Denmark	2014年7月12日
Current outcomes of donepezil treatment in Japan: a comparison between the Phase 3 trial and recent post-marketing (J-GOLD) studies	Alzheimer's Association International Conference 2014; Copenhagen, Denmark	2014年7月12日
Workaholism as a risk factor for depression and disabling back pain among Japanese workers	The 41st International Society for the Study of the Lumbar Spine; Seoul, Korea	2014年6月30日
Usefulness of 'One Stretch', a simple, daily, standing back extercise, for the prevention of onset of aggravation of low back pain in care workers	The 41st International Society for the Study of the Lumbar Spine; Seoul, Korea	2014年6月30日
Identification of risk factors for new-onset sciatica in Japanese workers: findings from the Japan epidemiological research of Occupation-related Back pain (JOB) study	The 23rd International Epidemiology in Occupational Health Conference; Utrecht, Netherlands	2013年6月18日
Potential risk factors of persistent low back pain developing from mild LBP in Japanese workers; findings from a prospective study	The 40th International Society for the Study of the Lumbar Spine; Scottsdale, US	2013年4月30日
言語妥当性を担保した日本語版Revised Fibromyalgia Impact Questionnaire (JFIQR) の作成	第57回日本リウマチ学会総会・学術集会	2013年4月18日

日本語版Tampa Scale for Kinesiophobia (TSK-J) の開発：言語的妥当性を担保した翻訳版の作成	第5回日本運動器疼痛学会	2012年11月17日
日本語版 Fear-Avoidance Beliefs Questionnaire (FABQ-J) の計量心理学的検討	第20回日本腰痛学会	2012年11月13日
Risk factors for new-onset of low back disability among Japanese workers; results from the Cultural and Psychosocial Influences on Disability (CUPID) study	The 22nd International Epidemiology in Occupational Health Conference; Oxford, UK	2011年8月29日
日本語版Neck Disability Indexの妥当性検証	第40回日本脊椎脊髄病学会	2011年4月22日
日本語版Fear-Avoidance Beliefs Questionnaire (FABQ) の開発：言語的妥当性を担保した翻訳版の作成	第18回日本腰痛学会	2010年10月31日
Risk factors for new onset of low back disability among Japanese workers; findings from the Japan epidemiological research of Occupation-related Back pain (JOB) study	The 21st International Epidemiology in Occupational Health Conference; Taipei, Taiwan	2010年4月21日
日本語版Zurich Claudication Questionnaire (ZCQ) の開発：言語的妥当性を担保した翻訳版の作成	第17回日本腰痛学会	2009年11月21日