

Examples of our experience in supporting clinical research or questionnaire development  
(published papers with our employee's names or our name in author, acknowledgment or text)

Title/Journal	Listed as/in
Characteristics of patients with longer treatment period of lenvatinib for unresectable hepatocellular carcinoma: a post-hoc analysis of post-marketing surveillance study in Japan. PLOS ONE. 2024; 19(3): e0298420.	Acknowledgment
Association between glycemic control and patient-reported outcomes in adults with type 1 diabetes in Japan: the SAGE study subanalysis. Diabetol Int. 2023; 15(2): 212-222.	Acknowledgment
Validation and comparison of the coding algorithms to identify people with migraine using Japanese claims data. Front Neurol. 2023; 14: 1231351.	Acknowledgment
Real-world treatment patterns of subsequent therapy after palbociclib in patients with advanced breast cancer in Japan. Breast. 2023; 70: 1-7.	Acknowledgment
Real-world treatment patterns of novel drugs in relapsed or refractory acute lymphoblastic leukemia patients in Japan. Future Oncol. 2023; 19(19): 1343-1356.	Acknowledgment
Exploring changes in attitudes, behaviors, and self-measured health data through lifestyle modification support by community pharmacists: Suito-Ogaki Selfcare (SOS) trial. Integr Pharm Res Pract. 2023; 12: 87-99.	Acknowledgment
Impact of tolvaptan add-on treatment on patients with heart failure requiring long-term congestion management: a retrospective cohort study using a medical claim database in Japan. J Cardiol. 2023; 82(1): 35-42.	Acknowledgment
Perioperative analgesics and anesthesia as risk factors for postoperative chronic opioid use in patients undergoing total knee arthroplasty: a retrospective cohort study using Japanese hospital claims data. Drugs - Real World Outcomes. 2023; 10(2): 331-340.	Author and Acknowledgment
Safety and effectiveness of molnupiravir (LAGEVRIO®) capsules in Japanese patients with COVID-19: interim report of post-marketing surveillance in Japan. Infect Dis Ther. 2023; 12(4): 1119-1136.	Acknowledgment
Validation of algorithms to identify colorectal cancer patients from administrative claims data of a Japanese hospital. BMC Health Serv Res. 2023; 23(1): 274.	Author and Acknowledgment
Linguistic and psychometric validation of the cognition bolt-on version of the Japanese EQ-5D-5L for the elderly. J Alzheimers Dis. 2023; 91(4): 1447-1458.	Author and Acknowledgment
Clinical characteristics, medication use, and impact of primary headache on daily activities: an observational study using linked online survey and medical claims data in Japan. BMC Neurol. 2023; 23(1): 80.	Acknowledgment
An attempt to replicate randomized trials of diabetes treatments using a Japanese administrative claims and health checkup database: a feasibility study. Drugs - Real World Outcomes. 2023; 10(2): 235-247.	Author and Acknowledgment
Validation of algorithms to identify bone metastases using administrative claims data in a Japanese hospital. Drugs - Real World Outcomes. 2023; 10(2) 187-194.	Author and Acknowledgment
Safety and effectiveness of lenvatinib in patients with unresectable hepatocellular carcinoma in real-world clinical practice: an observational post-marketing study in Japan. Drugs - Real World Outcomes. 2023; 10(2): 195-205.	Acknowledgment
Impact of "time zero" of follow-up settings in a comparative effectiveness study using real-world data with a non-user comparator: comparison of six different settings. Drugs - Real World Outcomes. 2023; 10(1): 107-117.	Author and Acknowledgment
Cardiometabolic risk reductions in patients with type 2 diabetes mellitus newly treated with a sodium–glucose cotransporter 2 inhibitor versus a dipeptidyl peptidase-4 inhibitor: A real-world administrative database study in Japan. J Diabetes Investig. 2023; 14(3): 404-416.	Author and Acknowledgment

Physician consultation rates and characteristics among workers with chronic pain or headache who participated in a behavioural change program: a retrospective database analysis using real-world healthcare data. <i>BMJ Open</i> . 2022; 12(11): e056846.	Acknowledgment
Effect of parenteral energy or amino acid doses on in-hospital mortality among patients with aspiration pneumonia: a cohort medical claims database study. <i>J Gerontol A Biol Sci Med Sci</i> . 2022; 77(8): 1683-1690.	Acknowledgment
Treatment discontinuation among patients with schizophrenia treated with brexpiprazole and other oral atypical antipsychotics in Japan: A retrospective observational study. <i>Adv Ther</i> . 2022; 39(9): 4299-4314.	Acknowledgment
Reduction in asthma exacerbation rate after mepolizumab treatment initiation in patients with severe asthma: A real-world database study in Japan. <i>Pulm Pharmacol Ther</i> . 2022; 75: 102130.	Acknowledgment
Effect of donepezil for dementia prevention in Parkinson's disease with severe hyposmia (The DASH-PD study): A randomized long-term placebo-controlled trial. <i>eClinicalMedicine</i> . 2022; 51: 101571.	Author
The use of a network analysis to identify associations and temporal patterns among non-communicable diseases in Japan based on a large medical claims database. <i>Drugs - Real World Outcomes</i> . 2022; 9(3): 463-476.	Author, Funding
A study to investigate the prevalence of headache disorders and migraine among people registered in a health insurance association in Japan. <i>J Headache Pain</i> . 2022; 23(1): 70.	Acknowledgment
Association between HbA1c levels and diabetic peripheral neuropathy: A case-control study of patients with type 2 diabetes using claims data. <i>Drugs - Real World Outcomes</i> . 2022; 9(3): 403-414.	Acknowledgment
Safety and effectiveness of glecaprevir/pibrentasvir in patients with chronic hepatitis C or compensated cirrhosis in daily clinical practice: a drug-use results survey. <i>Kanzo</i> . 2022; 63(3): 120-150.	Acknowledgment
Real-world treatment patterns of palbociclib and blood count monitoring in patients with advanced breast cancer in Japan. <i>Future Oncol</i> . 2022; 18(17): 2101-2111.	Acknowledgment
Comparative analysis of medical costs after hepatectomy versus radiofrequency ablation in patients with hepatocellular carcinoma in real-world clinical practice. <i>Hepatol Res</i> . 2022; 52(5): 471-478.	Acknowledgment
Pharmacotherapy for low back pain and sciatica: A narrative review. <i>J Spine Res</i> . 2022; 13(2): 67-86.	Acknowledgment
Investigation of the usefulness of EDC + ePRO + wearable device –A collaborative study–. <i>Prog Med</i> . 2022; 42(1): 91-100.	Acknowledgment
Internet survey on factors associated with care-seeking behaviours of people with chronic musculoskeletal pain in Japan. <i>J Orthop Surg (Hong Kong)</i> . 2021; 29(3): 23094990211044836.	Funding
A post marketing survey of intravenous daptomycin in Japanese patients. <i>J. New Rem. &amp; Clin</i> . 2021; 70: 1189-1215.	Acknowledgment
Nutritional management in inpatients with aspiration pneumonia: a cohort medical claims database study. <i>Arch Gerontol Geriatr</i> . 2021; 95: 104398.	Acknowledgment
Treatment patterns and medical costs after hepatectomy in real-world practice for patients with hepatocellular carcinoma in Japan. <i>Hepatol Res</i> . 2021; 51(10): 1073-1081.	Acknowledgment
Understanding confidence in the human papillomavirus vaccine in Japan: a web-based survey of mothers, female adolescents, and healthcare professionals. <i>Hum Vaccin Immunother</i> . 2021; 17(9): 3102-3112.	Acknowledgment
Real-world treatment patterns of patients with atopic dermatitis in Japan: analysis of the JMDC claims database. <i>J Cutan Immunol Allergy</i> . 2021; 00: 1–11.	Acknowledgment

Multiple comorbidities increase the risk of death from invasive pneumococcal disease under the age of 65 years. J Infect Chemother. 2021; 27(9): 1311-1318.	Acknowledgment
Glycemic control status, diabetes management patterns, and clinical characteristics of adults with type 1 diabetes in Japan: study of adults' glycemia in T1DM subanalysis. Diabetol Int. 2021; 12(4): 460-473.	Acknowledgment
Real-world treatment patterns and adherence to oral medication among patients with bipolar disorders: a retrospective, observational study using a healthcare claims database. Neuropsychiatr Dis Treat. 2021; 17: 821-833.	Acknowledgment
Real-world hospital mortality of liver cirrhosis inpatients in Japan: a large-scale cohort study using a medical claims database: prognosis of liver cirrhosis. Hepatol Res. 2021; 51(6): 682-693.	Acknowledgment
Effectiveness of pregabalin treatment for neuropathic pain in patients with spine diseases: a pooled analysis of two multicenter observational studies in Japan. J Pain Res. 2021; 14: 757-771.	Acknowledgment
Antidyslipidemic drug prescriptions and lipid control status after unfavorable annual health checkup results: a retrospective cohort study using a health insurance database. Drugs - Real World Outcomes. 2021; 8(2): 227-240.	Acknowledgment
Indices of peripheral leukocytes predict longer overall survival in breast cancer patients on eribulin in Japan. Breast Cancer. 2021; 28(4): 945-955.	Acknowledgment
Changes in the penetration rate of biosimilar infliximab within Japan using a Japanese claims database. Clinicoecon Outcomes Res. 2021; 13: 145-153.	Author
Relationship of annual health checkups with the subject's subsequent behavior of cardiovascular risk management in a real-world setting in Japan: a retrospective cohort study on changes in antihypertensive drug prescription and blood pressure from 2015 to 2017. Drugs Real World Outcomes. 2021; 8(2): 215-225.	Acknowledgment
Risk of hypoglycemia in Japanese people with type 2 diabetes mellitus who initiated or switched to insulin glargine 300 U/mL: a subgroup analysis of 12-month post-marketing surveillance study (X-STAR study). Diabetes Res Clin Pract. 2021; 172: 108647.	Acknowledgment
Clinical impact of prescribed doses of nutrients for patients exclusively receiving parenteral nutrition in Japanese hospitals: a retrospective cohort study. J Parenter Enteral Nutr. 2021; 45(7): 1514-1522.	Acknowledgment
Treatment and healthcare cost among patients with hip or knee osteoarthritis: a cross-sectional study using a real-world claims database in Japan between 2013 and 2019. Clin Drug Investig. 2020; 40(11): 1071-1084.	Author and Acknowledgment
Comparison of hospital length of stay of acute ischemic stroke patients with non-valvular atrial fibrillation started on rivaroxaban or warfarin treatment during hospitalization. J Med Econ. 2020; 23(12): 1379-1388.	Author and Acknowledgment
Real-world data on the use of insulin glargine 300 U/mL in Japanese patients with type 1 diabetes: twelve-month results from a post-marketing surveillance study (X-STAR study). Expert Opin Pharmacother. 2021; 22(2): 249-256.	Acknowledgment
Effectiveness and safety of insulin glargine 300 unit/mL in Japanese type 2 diabetes mellitus patients: a 12-month post-marketing surveillance study (X-STAR study). Expert Opin Pharmacother. 2020; 21: 1771-1780.	Acknowledgment
A survey on total parenteral nutrition in 55,000 hospitalized patients: retrospective cohort study using a medical claims database. Clin Nutr ESPEN. 2020; 39:198-205.	Author and Acknowledgment
Safety and effectiveness of lenvatinib in 594 patients with unresectable thyroid cancer in an all-case post-marketing observational study in Japan. Adv Ther. 2020; 37(9): 3850-3862.	Acknowledgment
Psychometric properties of the Japanese version of the Kansas City Cardiomyopathy Questionnaire in Japanese patients with chronic heart failure. Health Qual Life Outcomes. 2020; 18(1): 236.	Acknowledgment
Incidence of and risk factors for recurrent Clostridioidesdifficile infection in Japan using a claims database: a retrospective cohort study. Anaerobe. 2020; 61: 102139.	Acknowledgment

Prevalence of ascites and its treatment in inpatients with liver cirrhosis: a cohort study using a Japanese medical claims database. <i>Kanzo</i> . 2020; 61(6): 314-325.	Acknowledgment
Treatment persistence between long-acting injectable versus orally administered aripiprazole among patients with schizophrenia in a real-world clinical setting in Japan. <i>Adv Ther</i> . 2020; 37(7): 3324-3336.	Acknowledgment
Effectiveness and safety of eribulin in Japanese patients with HER2-negative, advanced breast cancer: a 2-year post-marketing observational study in a real-world setting. <i>Invest New Drugs</i> . 2020; 38(5): 1540-1549.	Acknowledgment
Long-term tolerability and effectiveness of raltegravir in Japanese patients: results from post-marketing surveillance. <i>PLoS One</i> . 2019; 14(1): e0210384.	Acknowledgment
Patterns of drug treatment in patients with osteoarthritis and chronic low back pain in Japan: a retrospective database study. <i>J Pain Res</i> . 2019; 12: 1631-1648.	Acknowledgment
Daily practice and prognostic factors for pneumonia caused by methicillin-resistant <i>Staphylococcus aureus</i> in Japan: a multicenter prospective observational cohort study. <i>J Infect Chemother</i> . 2020; 26(2): 242-251.	Acknowledgment
Clinical research on the effect of difference in properties of external preparations for skin on application amount: an exploratory investigation of the relationship between the base usability and application amount by questionnaire in adult subjects. <i>Yakugaku Zasshi</i> . 2019; 139(10): 1313-1325.	Acknowledgment
Development of a Japanese version of the Örebro Musculoskeletal Pain Screening Questionnaire (ÖMPSQ-J) and the short form version: translation and linguistic validation. <i>Journal of Musculoskeletal Pain Research</i> . 2019; 11: 35-48.	Author
A cost-effectiveness analysis of pregabalin for the treatment of patients with chronic cervical pain with a neuropathic component in Japan. <i>J Pain Res</i> . 2019; 12: 2785-2797.	Author
Safety and effectiveness of tofogliflozin in elderly Japanese patients with type 2 diabetes mellitus: a subanalysis of a post-marketing study (J-STEP/EL Study). <i>J Diabetes Investig</i> . 2020; 11(2): 405-416.	Acknowledgment
Sustained reduction in rotavirus-coded hospitalizations in children aged <5 years after introduction of self-financed rotavirus vaccines in Japan. <i>Hum Vaccin Immunother</i> . 2020; 16(1): 132-137.	Acknowledgment
Safety and effectiveness of tofogliflozin in Japanese patients with type 2 diabetes mellitus in real-world practice: Results of 12-month interim analysis of a long-term post-marketing surveillance study (J-STEP/LT). <i>J Diabetes Investig</i> . 2020; 11(1): 132-141.	Acknowledgment
Switch rates, time-to-switch, and switch patterns of antiretroviral therapy in people living with human immunodeficiency virus in Japan, in a hospital-claim database. <i>BMC Infect Dis</i> . 2019; 19(1): 505.	Acknowledgment
Pharmacotherapy decision-making among patients with breast cancer in Japan: results of an online survey. <i>Breast Cancer</i> . 2019; 26(6): 799-807.	Acknowledgment
Assessment of municipalities' sociodemographic characteristics and efforts associated with the high specific medical checkup rates. <i>Kosei no Shihyo (Journal of Health and Welfare Statistics)</i> . 2019; 66(4): 29-35.	Acknowledgment
Use of a healthcare claims database for post-marketing safety assessments of eribulin in Japan: a comparative assessment with a prospective post-marketing surveillance study. <i>Drugs Real World Outcomes</i> . 2019; 6(1): 27-35.	Author
Secular trend in the incidence of Japanese employees with thyroid cancer undergoing thyroidectomy from 2005 to 2014: a retrospective descriptive study using an employment-based insurance claims database. <i>J of thyroid cancer</i> . 2019; 1: 5-15.	-
Ultra-high-dose methylcobalamin in amyotrophic lateral sclerosis: a long-term phase II/III randomised controlled study. <i>J Neurol Neurosurg Psychiatry</i> . 2019; 90(4): 451-457.	Acknowledgment
Differences in the incidence of postoperative pneumonia after percutaneous endoscopic gastrostomy between liquid and semi-solid nutrient administration. <i>Eur J Clin Nutr</i> . 2019; 73(2): 250-257.	Acknowledgment and text (methods)

Assessment of risk factors for non-specific chronic disabling low back pain in Japanese workers—findings from the CUPID (Cultural and Psychosocial Influences on Disability) study. <i>Industrial Health</i> . 2019; 57(4): 503-510.	Author
Greater burden of chronic comorbidities and co-medications among people living with HIV versus people without HIV in Japan: a hospital claims database study. <i>J Infect Chemother</i> . 2019; 25(2): 89-95.	Acknowledgment
Chemotherapy-induced peripheral neuropathy in breast cancer patients treated with eribulin: interim data from a post-marketing observational study. <i>Breast Cancer</i> . 2019; 26(2): 235-243.	Acknowledgment
Pondering the “psychogenic pain”: proposal for using the term “cognitively perceived pain”. <i>Pain Research</i> . 2018; 33(3): 183-192.	Disclosure
Healthcare burden of recurrent <i>Clostridioides difficile</i> infection in Japan: a retrospective database study. <i>J Infect Chemother</i> . 2018; 24(11): 892-901.	Author
Recent trends in the practice of procedural sedation under local anesthesia for catheter ablation, gastrointestinal endoscopy, and endoscopic surgery in Japan: a retrospective database study in clinical practice from 2012 to 2015. <i>Drugs Real World Outcomes</i> . 2018; 5(3): 137-147.	Author
Low-density lipoprotein cholesterol goal attainment rates by initial statin monotherapy among patients with dyslipidemia and high cardiovascular risk in Japan—a retrospective database analysis. <i>Circ J</i> . 2018; 82(6): 1605-1613.	Acknowledgment
Comorbidities and the use of comedications in people living with HIV on antiretroviral therapy in Japan: a cross-sectional study using a hospital claims database. <i>BMJ Open</i> . 2018; 8(6): e019985.	Acknowledgment
Comorbidities and co-medications in populations with and without chronic hepatitis C virus infection in Japan between 2015 and 2016. <i>BMC Infect Dis</i> . 2018; 18(1): 237.	Acknowledgment
Post-marketing surveillance of donepezil in Japanese patients with Alzheimer's disease using ADAS-J cog: final reports and constructing a model for ADAS-J cog scores. <i>Japanese Journal of Geriatric Psychiatry</i> . 2018; 29(4): 413-426.	Text (methods)
Pregabalin as an analgesic option for patients undergoing thoracotomy: cost analysis of pregabalin versus epidural analgesia for post-thoracotomy pain relief. <i>J Thorac Dis</i> . 2018; 10(4): 2321-2330.	Author
Development of the Japanese Core Outcome Measures Index (COMI): cross-cultural adaptation and psychometric validation. <i>BMC Musculoskelet Disord</i> . 2018; 19(1): 71.	Author
Long-term safety and efficacy of tofogliflozin as add-on to insulin in patients with type 2 diabetes: results from a 52-week, multicentre, randomized, double-blind, open-label extension, Phase 4 study in Japan (J-STEP/INS). <i>Diabetes Obes Metab</i> . 2018; 20(5): 1176-1185.	Acknowledgment
Dipeptidyl peptidase-4 inhibitors as preferable oral hypoglycemic agents in terms of treatment satisfaction: results from a multicenter, 12-week, open label, randomized controlled study in Japan (PREFERENCE 4 study). <i>J Diabetes Investig</i> . 2018; 9(1): 137-145.	Acknowledgment
Decline of rotavirus-coded hospitalizations in children under 5 years: a report from Japan where rotavirus vaccines are self-financed. <i>Vaccine</i> . 2018; 36(19): 2727-2732.	Acknowledgment
Clinical and economic burdens experienced by patients with painful diabetic peripheral neuropathy: an observational study using a Japanese claims database. <i>PLoS One</i> . 2017; 12(10): e0187250.	Author
Assessment of potential risk factors for new onset disabling low back pain in Japanese workers: findings from the CUPID (cultural and psychosocial influences on disability) study. <i>BMC Musculoskelet Disord</i> . 2017; 18(1): 334.	Author
Semi-quantitative assessment of tumor spread through air spaces (STAS) in early stage lung adenocarcinomas. <i>J Thorac Oncol</i> . 2017; 12(7): 1046-1051.	Acknowledgment and text (methods)
Development of a Japanese version of the Somatic Symptom Scale-8: psychometric validity and internal consistency. <i>General Hospital Psychiatry</i> . 2017; 45: 7-11.	Author

Development of the Japanese version of the leeds assessment of the neuropathic symptoms and signs pain scale: diagnostic utility in a clinical setting. <i>Pain Pract.</i> 2017; 17(6): 800-807.	Author
The Japanese version of the STarT Back Tool predicts 6-month clinical outcomes of low back pain. <i>J Orthop Sci.</i> 2017; 22(2): 224-229.	Author
Usefulness of a simple self-administered joint condition assessment sheet to predict the need for orthopaedic intervention in the management of haemophilic arthropathy. <i>Haemophilia.</i> 2017; 23(2): e116-e123.	Acknowledgment
Psychometric assessment of the Japanese version of the revised fibromyalgia impact questionnaire: reliability and validity. <i>Int J Rheum Dis.</i> 2017; 20(9): 1088-1094.	Author
Comparison of visceral adipose tissue area measured by computed tomography at the different vertebrate level—sub-analysis using data of Japanese patients derived from International Study of the Prediction of Intra-Abdominal Adiposity and its Relationship with CardioMetabolic Risk/Intra-Abdominal Adiposity (INSPIRE-ME-IAA)—. <i>J Clin Therap Med.</i> 2016; 32(9): 783-791.	Acknowledgment
Distribution of ankle-brachial index among inpatients with cardiovascular disease: analysis using the Kumamoto University Hospital medical database. <i>Annals of vascular diseases,</i> 2016; 9(1): 22-29.	Acknowledgment
Impact of patient attitudes and beliefs to insulin therapy upon initiation, and their attitudinal changes after initiation: the DAWN Japan study. <i>Curr Med Res Opin,</i> 2016; 32(4): 681-686.	Acknowledgment
Host factors and biomarkers associated with poor outcomes in adults with invasive pneumococcal disease. <i>PLoS ONE.</i> 2016; 11(1): e0147877.	Acknowledgment
Pretreatment cognitive profile likely to benefit from donepezil treatment in dementia with lewy bodies: pooled analyses of two randomized controlled trials. <i>Dement Geriatr Cogn Disord.</i> 2016; 42: 58-68.	Acknowledgment
Development of a linguistically validated Japanese version of the Somatic Symptom Scale-8 (SSS-8). <i>Japanese Journal of Psychosomatic Medicine (Tokyo).</i> 2016; 56(9): 931-937.	Author
Liver-related safety assessment of green tea extracts in humans: a systematic review of randomized controlled trials. <i>EJCN.</i> 2016; 70(11): 1221-1229.	Author
Psychometric assessment of the Japanese version of the Zurich Claudication Questionnaire (ZCQ): reliability and validity. <i>PLoS ONE.</i> 2016; 11(7): e0160183.	Author
Low back pain prevention exercise – My method for workers –. <i>MB Med Reha.</i> 2016; No.198: 63-69. (in Japanese)	Author
Psychometric properties of the Japanese version of the STarT Back Tool in patients with low back pain. <i>PLoS ONE.</i> 2016; 11(3): e0152019.	Author
Potential risk factors for onset of severe neck and shoulder discomfort (Katakori) in urban Japanese workers. <i>Ind Health.</i> 2016; 54(3): 230-236.	Author
Efficacy and safety assessment of basal supported oral therapy (BOT) with insulin glargine in a real-life clinical setting, stratified by concomitant orally administered antidiabetic agent (OAD) regimens including dipeptidyl peptidase-4 inhibitor (DPP-4i): subanalysis of the ALOHA2 study, drug-use surveillance in Japan. <i>Diabetol Int.</i> 2016; 7(3): 299-307.	Acknowledgment
Predictors for achieving target glycemic control in Japanese patients with type 2 diabetes after initiation of basal supported oral therapy using insulin glargine: sub-analysis of the ALOHA2 study, drug use surveillance in Japan. <i>Diabetol Int.</i> 2016; 7(2): 188-198.	Acknowledgment
Japanese postmarketing surveillance of clopidogrel in patients with non-ST-segment elevation acute coronary syndrome, stable angina, old myocardial infarction, and ST-segment elevation myocardial infarction after percutaneous coronary intervention in a real-life setting: the final report (J-PLACE Final). <i>Cardiovasc Interv Ther.</i> 2016; 31(2): 101-113.	Acknowledgment
The relationship between patient perception of healthcare provision by professionals and the self-care activity of patients with diabetes: Japanese subgroup analysis of the second Diabetes Attitudes, Wishes, and Needs (DAWN2) study. <i>Diabetol Int.</i> 2016; 7(2): 111-118.	Acknowledgment

Disease state changes and safety of long-term donepezil hydrochloride administration in patients with Alzheimer's disease: interim results from the long-term, large-scale J-GOLD study in Japan. <i>Psychogeriatrics</i> . 2016; 2: 107-115.	Author and text (methods)
Development of the Japanese version of the Menorrhagia Multi-Attribute Scale(MMAS): translation and linguistic validation. <i>Sanka to Fujinka</i> . 2015; 82(11): 1299-1305. (in Japanese)	Author
Psychometric properties of the Japanese version of the Tampa Scale for Kinesiophobia (TSK-J) in patients with whiplash neck injury pain and/or low back pain. <i>J Orthop Sci</i> . 2015; 20(6): 985-992.	Author
Can standing back extension exercise improve or prevent low back pain in Japanese care workers? <i>J Man Manip Ther</i> . 2015; 23(4): 205-209.	Author
Adequacy of using consensus guidelines for diagnosis of dementia with lewy bodies in clinical trials for drug development. <i>Dement Geriatr Cogn Disord</i> . 2015; 41(1-2): 55-67.	Acknowledgment
Assessment of psychosocial risk factors for the development of non-specific chronic disabling low back pain in Japanese workers—findings from the Japan epidemiological research of Occupation-related Back pain (JOB) study. <i>Ind Health</i> . 2015; 53(4): 368-377.	Author
Effects of donepezil on extrapyramidal symptoms in patients with dementia with Lewy bodies: a secondary pooled analysis of two randomized-controlled and two open-label long-term extension studies. <i>Dement Geriatr Cogn Disord</i> . 2015; 40(3-4): 186-198.	Acknowledgment
Improved treatment satisfaction and self-reported health status after introduction of basal-supported oral therapy using insulin glargine in patients with type 2 diabetes: sub-analysis of ALOHA2 Study. <i>Diabetes Ther</i> . 2015; 6(2): 153-171.	Acknowledgment
The effects of bazedoxifene on bone, glucose, and lipid metabolism in postmenopausal women with type 2 diabetes: an exploratory pilot study. <i>J Clin Med Res</i> . 2015; 7(10): 762-769.	Acknowledgment
Effect of postoperative administration of pregabalin for post-thoracotomy pain: a randomized study. <i>J Cardiothorac Vasc Anesth</i> . 2015; 29(6): 1567-1572.	Acknowledgment and text (methods)
Epidemiology resesrach on low back pain in Japanese workers. <i>JJOMT</i> . 2015; 63(6): 329-336.	Author
Current situation and issues in titration of basal insulin. <i>Therapeutic Research</i> . 2015; 36(5): 403-411.	Acknowledgment
Linguistic validation of the English version of the Core Lower Urinary Tract Symptom Score. <i>Int J Urol</i> . 2014; 21(12):1291-1292.	Text (methods)
Development of the Japanese version of the Leeds Assessment of Neuropathic Symptoms and Signs Pain Scale: translation and linguistic validation. <i>Pain Clinic</i> . 2014; 35(7): 933-940. (in Japanese)	Author
Psychometric validation of the English version of the overactive bladder symptom score. <i>Urology</i> . 2014; 84(1): 46-50.	Text (methods)
Development of the Japanese version of the Revised Fibromyalgia Impact Questionnaire (JFIQR): translation and linguistic validation. <i>Clinical Rheumatology and Related Research</i> . 2014; 26(1): 35-44. (in Japanese)	Author
Development of the Japanese version of the London Fibromyalgia Epidemiology Study Screening Questionnaire (LFESSQ): translation and linguistic validation. <i>Clinical Rheumatology and Related Research</i> . 2014; 26(2): 130-136. (in Japanese)	Author
Psychometric properties of the Japanese version of the Fear-Avoidance Beliefs Questionnaire (FABQ). <i>J Orthop Sci</i> . 2014; 19(1): 26-32.	Author
Linguistic validation of the English version of the Overactive Bladder Symptom Score. <i>Int J Urol</i> . 2014; 21(2): 229.	Text

Central nervous system effects of the second-generation antihistamines marketed in Japan—review of inter-drug differences using the Proportional Impairment Ratio (PIR)—. PLoS ONE. 2014; 9(12): e114336.	Author
Gaps between actual and desired sex life: web survey of 5,665 Japanese women. J Sex Marital Ther. 2014; 40(1): 33-42.	Text (methods)
Potential risk factors of persistent low back pain developing from mild low back pain in urban Japanese workers. PLoS ONE. 2014; 9(4): e93924.	Author
Validity, reliability and responsiveness of the Japanese version of the Neck Disability Index. J Orthop Sci, 2013; 18(1); 14-21.	Author
Development of a Japanese version of the Tampa Scale for Kinesiophobia (TSK-J) : translation and linguistic validation. Rinshou Seikeigeka. 2013; 48(1): 13-19. (in Japanese)	Author
Development of the Japanese version of the STarT (Subgrouping for Targeted Treatment) Back Screening Tool; translation and linguistic validation. Journal of Musculoskeletal Pain Research. 2013; 5(1): 11-19. (in Japanese)	Author
Identification of risk factors for new-onset sciatica in Japanese workers: findings from the Japan epidemiological research of Occupation-related Back pain study. Spine (Phila Pa 1976). 2013; 38(26): E1691-E1700.	Author
Maintenance therapy with pegaptanib sodium for neovascular age-related macular degeneration: an exploratory study in Japanese patients (LEVEL-J study). Jpn J Ophthalmol. 2013; 57(5): 417-423.	Acknowledgment
Development and validation of a new questionnaire assessing quality of life in adults with hypopituitarism: Adult Hypopituitarism Questionnaire (AHQ). PLoS ONE. 2012; 7(9): e44304.	Text (methods)
Development of the Japanese version of the brief symptom inventory-somatization scale: translation and linguistic validation. Seikei Geka. 2012; 63(2): 149-153. (in Japanese)	Author
Potential risk factors for new onset of back pain disability in Japanese workers: findings from the Japan epidemiological research of occupation-related back pain study. Spine (Phila Pa 1976). 2012; 37(15): 1324-1333.	Author
Recombinant activated factor VII (rFVIIa/NovoSeven®) in the management of severe postpartum haemorrhage: initial report of a multicentre case series in Japan. Int J Hematol. 2012; 95(1): 57-63.	Acknowledgment
Development of the Japanese version of the Fear-Avoidance Beliefs Questionnaire (FABQ-J): translation and linguistic validation. Seikei Geka Gakkai. 2011; 62(12): 1301-1306. (in Japanese)	Author
Responsiveness and minimal clinically important change in overactive bladder symptom score. Urology. 2011; 78(4): 768-773.	Acknowledgment
Development of the Japanese version of the Fibromyalgia Impact Questionnaire (JFIQ): psychometric assessments of reliability and validity. Int J Rheum Dis. 2011; 14(1): 74-80.	Author and text (methods)
Comparison of physician's advice for non-specific acute low back pain in Japanese workers: advice to rest versus advice to stay active. Ind Health. 2011; 49(2): 203-208.	Author
Solifenacin as add-on therapy for overactive bladder symptoms in men treated for lower urinary tract symptoms—ASSIST, randomized controlled study. Urology. 2011; 78(1): 126-133.	Acknowledgment
Development of the Japanese version of the Zurich claudication questionnaire (ZCQ): translation and linguistic validation. Nihon Seikei Geka Gakkai. 2010; 61(2): 159-165. (in Japanese)	Author
Development of the Japanese version of the Fibromyalgia Impact Questionnaire (JFIQ): translation and linguistic validation. Clin Rheumatol. 2008; 20: 19-28. (in Japanese)	Author
Opinion survey of women who have male sexual partners on 'better sex'. The Japanese journal of Impotence Research. 2007; 22(3): 287-300. (in Japanese)	Text (methods)
Psychometric properties of the Japanese version of the overactive bladder questionnaire (OAB-q). Journal of the Japan Neurogenic Bladder Society. 2006; 17(2): 250-256. (in Japanese)	Text (methods)



Examples of our experience in supporting conference presentations  
(oral or poster)

Title	Conference	Date
Impact of primary headache on daily activities from real world data in Japan	IHC 2023 - International Headache Congress; Seoul, Korea	September 14, 2023
Linguistic and psychometric validation of the "cognition" bolt-on version of the Japanese EQ-5D-5L (EQ-5D-5L+C) by adding a cognitive dimension for the elderly	ISPOR Europe 2022; Vienna, Austria	November 7, 2022
Safety and efficacy of long-term treatment with perampanel in Japanese patients with partial-onset seizure or primary generalized tonic-clonic seizure in a real-world setting	American Epilepsy Society Annual Meeting 2019; Baltimore, US	December 9, 2019
Patterns of drug treatments among patients with osteoarthritis and chronic low back pain and characteristics of patients with osteoarthritis who underwent any surgery related to osteoarthritis (in Japanese)	The 12th Japanese Association for the Study of Musculoskeletal Pain; Tokyo, Japan	November 30, 2019
Examination of the effectiveness and safety of insulin glargine 300 U/mL hospitalization introduction in insulin-naive patients with type 2 diabetes: X-STAR study sub-analysis (in Japanese)	The 62nd Annual Meeting of the Japan Diabetes Society; Sendai, Japan	May 23, 2019
Eribulin as first- or second-line chemotherapy for advanced or metastatic HER2-negative breast cancer: a real-world prospective study	The European Society for Medical Oncology 2018 Congress; Munich, Germany	October 19, 2018
The usability of commercial healthcare claims data in Japanese pharmacovigilance: a comparison with a prospective postmarketing study	The 34th International Conference on Pharmacoepidemiology & Therapeutic Risk Management; Prague, Czech republic	August 22, 2018
Treatment patterns and low-density lipoprotein cholesterol goal attainment rates among patients with dyslipidemia and high cardiovascular risk in Japan: a retrospective database analysis	The 82nd Annual Scientific Meeting of the Japanese Circulation Society; Osaka, Japan	March 25, 2018
A real-world observational study of eribulin in patients with soft-tissue sarcoma in Japan	The 22nd Annual Meeting of Connective Tissue Oncology Society; Hawaii, US	November 8, 2017
Prospective observational study of peripheral neuropathy in breast cancer patients treated with eribulin	The European Society for Medical Oncology 2017 Congress; Madrid, Spain	September 11, 2017
Diagnostic capability of "Spine paindetect" to screen for neuropathic pain caused by spinal disorders	The 44th Annual Meeting of the International Society for the Study of the Lumbar Spine; Athens, Greece	June 1, 2017
Effects of donepezil on the prognosis of Parkinson's disease with severe hyposmia—a multicenter randomized double-blind placebo-controlled trial (DASH-PD study)	The 20th International Congress of Parkinson's Disease and Movement Disorders; Berlin, Germany	June 22, 2016
Potential risk factors for onset of severe neck and shoulder discomfort (Katakori) in urban Japanese workers	The 89th Annual Meeting of Japan Society for Occupational Health; Fukushima, Japan	May 26, 2016
Psychometric validation of the Japanese version of the Somatic Symptom Scale-8 (SSS-8) (in Japanese)	The 8th Annual Meeting of Japanese Association for the Study of Musculoskeletal Pain; Nagoya, Japan	December 13, 2015
Association of risk factors for severe low back pain in workers in nursing homes (in Japanese)	The 23rd Annual Meeting of The Japanese Society of Lumbar Spine Disorders; Tokyo, Japan	November 15, 2015
The use of anti-epileptic drugs in Japanese patients with Alzheimer's disease on donepezil: results of large, real-world datasets (in Japanese)	The 49th Congress of The Japan Epilepsy Society; Nagasaki, Japan	October 30, 2015
Reality of treatment for patients with Alzheimer's disease in rapidly aging Japanese society: analysis using PMS data of donepezil (in Japanese)	The 34th Annual Meeting of Japan Society for Dementia Research; Aomori, Japan	October 2, 2015

Analysis of associations between high density lipoprotein structure and cholesterol efflux capacity in dyslipidemic patients treated with pitavastatin (High-Draw study) (in Japanese)	The 47th Annual Meeting of Japan Atherosclerosis Society; Sendai, Japan	July 9, 2015
Gender difference in the progression of Alzheimer's disease in Japanese patients treated with donepezil: results of the J-GOLD study.	Alzheimer's Association International Conference 2015; Washington, US	July 18, 2015
Risk factors for chronic non-specific low back pain in Japanese workers—JOB Study (in Japanese)	The 88th Annual Meeting of Japan Society for Occupational Health; Osaka, Japan	May 15, 2015
The potential for liver damage induced by green tea intake: a systematic review of clinical trials	The 25th Annual Scientific Meeting of the Japanese Epidemiology Association; Nagoya, Japan	January 23, 2015
Epidemiological study of low back pain in Japanese workers (in Japanese)	The 62nd Annual Meeting of Japanese Society of Occupational Medicine and Traumatology symposium 3 "Workers' management of low back pain"; Kobe, Japan	November 16, 2014
Psychometric properties of the Japanese version of the STarT Back screening tool (in Japanese)	The 22nd Annual Meeting of The Japanese Society of Lumbar Spine Disorders; Chiba, Japan	November 15, 2014
Potential predictive factors for cognitive response in patients with donepezil treatment: results from the J-GOLD study in Japan	Alzheimer's Association International Conference 2014; Copenhagen, Denmark	July 12, 2014
Impact of early donepezil treatment of Alzheimer's disease in a rapidly aging society: results from the J-GOLD study	Alzheimer's Association International Conference 2014; Copenhagen, Denmark	July 12, 2014
Current outcomes of donepezil treatment in Japan: a comparison between the Phase 3 trial and recent post-marketing (J-GOLD) studies	Alzheimer's Association International Conference 2014; Copenhagen, Denmark	July 12, 2014
Workaholism as a risk factor for depression and disabling back pain among Japanese workers	The 41st International Society for the Study of the Lumbar Spine; Seoul, Korea	June 30, 2014
Usefulness of 'One Stretch', a simple, daily, standing back extercise, for the prevention of onset of aggravation of low back pain in care workers	The 41st International Society for the Study of the Lumbar Spine; Seoul, Korea	June 30, 2014
Identification of risk factors for new-onset sciatica in Japanese workers: findings from the Japan epidemiological research of Occupation-related Back pain (JOB) study	The 23rd International Epidemiology in Occupational Health Conference; Utrecht, Netherlands	June 18, 2013
Potential risk factors of persistent low back pain developing from mild LBP in Japanese workers; findings from a prospective study	The 40th International Society for the Study of the Lumbar Spine; Scottsdale, US	April 30, 2013
Development of the Japanese version of the revised Fibromyalgia Impact Questionnaire (JFIQR): translation and linguistic validation (in Japanese)	The 57th Annual General Assembly and Scientific Meeting of the Japan College of Rheumatology; Kyoto, Japan	April 18, 2013
Development of the Japanese version of the Tampa Scale for Kinesiophobia (TSK-J): translation and linguistic validation (in Japanese)	The 5th Annual Meeting of the Japanese Association for the Study of Musculoskeletal Pain; Tokyo, Japan	November 17, 2012
Psychometric validation of the Japanese version of the Fear-Avoidance Beliefs Questionnaire (FABQ-J) (in Japanese)	The 20th Annual Meeting of The Japanese Society of Lumbar Spine Disorders; Kobe, Japan	November 3, 2012
Risk factors for new-onset of low back disability among Japanese workers; results from the Cultural and Psychosocial Influences on Disability (CUPID) study	The 22nd International Epidemiology in Occupational Health Conference; Oxford, UK	August 29, 2011
Psychometric validation of the Japanese version of the Neck Disability Index (in Japanese)	The 40th Annual Meeting of the Japanese Society for Spine Surgery and Related Research; Tokyo, Japan	April 22, 2011
Development of the Japanese version of the Fear-Avoidance Beliefs Questionnaire (FABQ): translation and linguistic validation (in Japanese)	The 18th Annual Meeting of the Japanese Society of Lumbar Spine Disorders; Hokkaido, Japan	October 31, 2010
Risk factors for new onset of low back disability among Japanese workers; findings from the Japan epidemiological research of Occupation-related Back pain (JOB) study	The 21st International Epidemiology in Occupational Health Conference; Taipei, Taiwan	April 21, 2010
Development of the Japanese version of the Zurich Claudication Questionnaire (ZCQ): translation and linguistic validation (in Japanese)	The 17th Annual Meeting of the Japanese Society of Lumbar Spine Disorders; Tokyo, Japan	November 21, 2009